

Supplementary appendix 6

This translation in Igbo was submitted by the authors and we reproduce it as supplied. It has not been peer reviewed. The Lancet's editorial processes have only been applied to the original in English, which should serve as reference for this manuscript.

Supplement to: Igboh LS, Roguski K, Marcenac P, et al. Timing of seasonal influenza epidemics for 25 countries in Africa during 2010–19: a retrospective analysis.
Lancet Glob Health 2023; **11**: e729–39.

Umi edemede ihe odide

Mgbakwasa-ukwu Iji data onyunyo obodo kpomkwem wee kowaa omume ntiwapu ɔrịa influenza nwere ike inye aka na ime mkpebi banyere oge igba ɔgwụ mgbochi maka ɔrịa influenza Anyị nyochara data onyunyo sitere na mba Afrika iji mara oge ntiwapu ɔrịa influenza iji nye aka n'imebe usoro igba ɔgwụ mgbochi ɔrịa na mba. Anyị ji data sentinel dì n'ihi ọha nke sitere na mba ndị na-eme mkpesa na netwok nke Global Influenza Surveillance and Response (GISRS) nke nwere data afọ 3-10 nke anakotara n'oge 2010-2019.

Usoro Anyị gbakotara izu-uka 3 ziri ezi na-agaghari agaghari nke nha nke ihe nlele dì mma maka nje influenza ma nyochaa oge ntiwapu ɔrịa na-efe efe site na iji usoro nchikota mkpokota.. Mbido na njedebe kwa oge ntiwapu ɔrịa na-efe efe bụ ihe akowara dika izu-uka mbu n'oge nha gafere ma ọ bụ gbadara n'uzo dì ala kwa afọ, n'otu n'otu, maka ȝizu-uka 3 nke na-esochi. Anyị kewaputara mba n'ime usoro ntiwapu ɔrịa na-efe efe ise: Ntiwapu ɔrịa na-efe efe nke Northern Hemisphere (NH) na-eme n'ọnwa Oktoba-Mach, Southern Hemisphere (SH) na-eme n'ọnwa Eprel-Septemba, n'uzo bù isi NH nwere ụfodụ ntiwapu ɔrịa na-efe efe n'ime ọnwa SH, n'uzo bù isi SH na ụfodụ ntiwapu ɔrịa na-efe efe n'ime ọnwa NH, na mbufe nke ɔrịa influenza n'ime afọ na-enweghi ihe a-na-ahụ anya nke NH ma ọ bụ pütara ihe nke SH.

Nchoputagasi Iri abụo na ise (74%) nke mba 34 GISRS nwere data karịrị ma ọ bụ dì nhata afọ 3, na anochiteanya 46% nke mba dì na Afrika, yana 89% nke ọnụ ögugụ mmadụ na Afrika. Enwere ntiwapu ɔrịa na-efe efe 248 n'ofe data afọ 236 nke obodo (nke etiti: 10; ögabata: 7-10 kwa mba). Mba isii (24%) nwere ükpurụ NH (Algeria, Egypt, Tunisia, Burkina Faso, Niger, na Morocco). Asatọ (32%) nwere ükpurụ nke NH n'uzo bù isi ma nwee ụfodụ ntiwapu ɔrịa na-efe efe nke SH (Cameroon, Ethiopia, Mali, Nigeria, Senegal, Togo, Tanzania, na Mozambique). Atọ (12%) nwere ükpurụ nke SH n'uzo bù isi ma nwee ụfodụ ntiwapu ɔrịa na-efe efe nke NH (Ghana, Kenya, na Uganda). Atọ (12%) nwere ukpuru nke SH (Central African Republic, Zambia, na South Africa). Ise (20%) enweghi NH ma ọ bụ mpụta ihe nke SH (Cote D'Ivoire, Democratic Republic of Congo, Rwanda, Madagascar, na Mauritius).

Nkowa Ọtụtu mba nwere oge ntiwapu ɔrịa na-efe efe nke ɔrịa influenza amatara maka iji gwa ndị ọchichị maka omume ɔrịa influenza nke na-abughị n'oge yana nke bụ n'oge mbufe ya, na-eduzi oge igba ɔgwụ mgbochi ɔrịa, ma kwalite mmemme ntinye aka n'oge.

Nkwado ego Ọnweghi