

III ROUND DELPHI							
COLOR CODE CLASS	SURGICAL DISEASE	Likert scale 4 (near totally agree)	%	Likert scale 5 (totally agree)	%	Reached Consensus Agreement (>=75%)	Decision
<b>RED class-Immediate - unstable patient</b>	Gynecological bleeding with unstable patient	6/43	<b>14</b>	35/43	<b>81.4</b>	Y	<b>Included</b>
<b>ORANGE class-Within 1 hour from diagnosis-</b> but as soon as possible, patient stable after target resuscitation but at risk of MOF	Acute bleeding non amenable of angio-embolization	16/43	<b>37.2</b>	24/43	<b>55.8</b>	Y	<b>Included</b>
	Ovarian torsion with unstable patient	19/43	<b>44.2</b>	13/43	<b>30.2</b>	N	IV round
	Strangulated (groin, incisional, ventral) hernia	12/43	<b>27.9</b>	18/43	<b>41.9</b>	N	IV round
	Gynecological bleeding in stable patient after resuscitation	19/43	<b>44.2</b>	10/43	<b>23.3</b>	N	IV round
	Ruptured tubo-ovarian abscess with septic shock	16/43	<b>37.2</b>	15/43	<b>34.9</b>	N	IV round

	<b>Complicated appendicitis (necrotic, appendicular abscess, perforated, presence of fecolith) associated with local/generalized peritonitis and signs of septic shock</b>	12/43	<b>27.9</b>	21/43	<b>48.8</b>	Y	<b>Included</b>
	<b>Complicated diverticulitis with local/generalized peritonitis and signs of septic shock (Hinchey III-IV)</b>	11/43	<b>25.6</b>	24/43	<b>55.8</b>	Y	<b>Included</b>
	<b>Perianal abscess with septic shock</b>	12/43	<b>27.9</b>	21/43	<b>48.8</b>	Y	<b>Included</b>
	Infected necrotizing hemorrhagic pancreatitis with signs of septic shock	11/43	<b>25.6</b>	17/43	<b>39.5</b>	N	
	Subdural hematoma	15/43	<b>34.9</b>	15/43	<b>34.9</b>	N	Cancelled
<b>YELLOW class-Within 6 hours from diagnosis-</b> stable patient with signs of sepsis	Anastomotic intestinal fistula/insufficiency in patient presenting sepsis signs	15/43	<b>34.9</b>	16/43	<b>37.2</b>	N	Modified definition; IV round
	Increasing intracranial pressure conditions	12/43	<b>27.9</b>	19/43	<b>44.2</b>	N	Cancelled
	<b>Complicated diverticulitis with sepsis (Hinchey III-IV)</b>	15/43	<b>34.9</b>	23/43	<b>53.5</b>	Y	<b>Included</b>

	<b>Complicated appendicitis (necrotic, appendicular abscess, perforated, presence of fecolith) associated with sepsis</b>	16/43	<b>37.2</b>	21/43	<b>48.8</b>	Y	<b>Included</b>
	Angioembolization of post splenic trauma pseudoaneurysm	10/43	<b>23.3</b>	14/43	<b>32.6</b>	N	Cancelled
	Foreign body with GUT obstruction (including endoscopy)	7/43	<b>16.3</b>	19/43	<b>44.2</b>	N	Cancelled
	Intraperitoneal bladder rupture	11/43	<b>25.6</b>	18/43	<b>41.9</b>	N	Cancelled
	Kidney transplant	8/43	<b>18.6</b>	12/43	<b>27.9</b>	N	Cancelled
	<b>Perianal abscess with sepsis</b>	23/43	<b>53.5</b>	13/43	<b>30.2</b>	Y	<b>Included</b>
	<b>Urolithiasis with sepsis</b>	21/43	<b>48.8</b>	13/43	<b>30.2</b>	Y	<b>Included</b>
	Urolithiasis with high risk of kidney injury	12/43	<b>27.9</b>	19/43	<b>44.2</b>	N	Modified definition; IV round

<b>GREEN class-Within 12 hours from diagnosis-stable patient with moderate risk of presenting hemodynamic instability and organs failure; admitted in surgical department for prompt medical treatment and monitoring</b>	Incarcerated abdominal wall hernia with intestinal obstruction	14/43	<b>32.6</b>	13/43	<b>30.2</b>	N	Modified; IV round
	Bilateral hydronephrosis	9/43	<b>20.9</b>	13/43	<b>30.2</b>	N	Cancelled
<b>BLUE class-Within 24 or 48 hour from diagnosis; stable patient with low risk of presenting organs failure, admitted in surgical department for clinical monitoring and medical treatment</b>	Bowel/intestinal obstruction after medical treatment failure	12/43	<b>27.9</b>	18/43	<b>41.9</b>	N	modified; IV round
	Perianal abscess	15/43	<b>34.9</b>	16/43	<b>37.2</b>	N	Modified; IV round
	<b>Cholecystitis</b>	10/43	<b>23.3</b>	24/43	<b>55.8</b>	<b>Y</b>	<b>Included</b>
	Urinary fistula	5/43	<b>11.6</b>	16/43	<b>37.2</b>	N	Cancelled
	Foreign body without intestinal obstruction	7/43	<b>16.3</b>	23/43	<b>53.5</b>	N	Cancelled
	Amputation due to osteomyelitis	14/43	<b>32.6</b>	16/43	<b>37.2</b>	N	Cancelled

	Acute appendicitis without peritoneal fluid	6/43	<b>14</b>	22/43	<b>51.2</b>	N	Modified; IV round
	A-V fistula for hemodialysis	10/43	<b>23.3</b>	12/43	<b>27.9</b>	N	Cancelled
	Complicated bones fractures (displaced, complex)	13/43	<b>30.2</b>	15/43	<b>34.9</b>	N	Cancelled
	Maxillo facial fractures	8/43	<b>18.6</b>	17/43	<b>39.5</b>	N	Cancelled