

eSupplement 2. Prescriptions, imaging, and interventions in the evaluation and treatment of 740 ocular and adnexal injuries

Intervention	No. (%)
Medication	
Antibiotic (topical)	338 (45.7%)
Cycloplegic	94 (12.7%)
Steroid (topical)	75 (10.1%)
Antibiotic (oral)	62 (8.4%)
Anti-histamine (oral)	57 (7.7%)
Opioid	34 (4.6%)
Ocular anti-hypertensive	11 (1.5%)
Other ^a	22 (2.9%)
Imaging studies	
Head CT	53 (7.2%)
B-scan ultrasonography	14 (1.9%)
X-ray	11 (1.5%)
Head MRI	2 (0.3%)
OCT	1 (0.1%)
Procedure	
Eye irrigation	43 (5.8%)
Foreign body removal (superficial)	16 (2.2%)
Eyelid/Adnexal laceration repair	14 (1.9%)
Foreign body removal (implanted)	5 (0.7%)
Exam under anesthesia	3 (0.4%)
Fracture repair	2 (0.3%)
Retina laser	1 (0.1%)
Corneal laceration repair	1 (0.1%)
Open globe repair	

^aIncludes topical anesthetics, IV antibiotics, topical antihistamines, and oral steroids