

Supplementary Table 2. Survey response distribution of bronchiectasis patients according to COVID-19 history by sex

		All (n=123)	COVID-19 (-) (n=99)	COVID-19 (+) (n=24)	<i>p</i> value
MARS, total mean score					
Female		23.2 ± 3.0	23.3 ± 3.0	22.8 ± 2.9	0.563 ^a
Male		21.6 ± 6.0	21.0 ± 6.2	23.1 ± 5.4	0.060 ^a
MARS, total score					
Female	≤20	8 (10.5)	6 (9.1)	2 (20)	0.282 ^c
	>20	68 (89.5)	60 (90.9)	8 (80)	
Male	≤20	9 (19.1)	8 (24.2)	1 (7.1)	0.173 ^b
	>20	38 (80.9)	25 (75.8)	13 (92.9)	
Beck Anxiety Scale					
Female	0-7 point(s)	33 (43.4)	29 (43.9)	4 (40)	0.185 ^c
	Mild	19 (25)	14 (21.2)	5 (50)	
	Moderate	14 (18.4)	14 (21.2)	0 (0)	
	Severe	10 (13.2)	9 (13.6)	1 (10)	
Male	0-7 point(s)	29 (61.7)	21 (63.6)	8 (57.1)	0.654 ^c
	Mild	12 (25.5)	7 (21.2)	5 (35.7)	
	Moderate	3 (6.4)	3 (9.1)	0 (0)	
	Severe	3 (6.4)	2 (6.1)	1 (7.1)	
IES-R total mean score					
Female		15.6 ± 18.1	16.3 ± 18.5	11.0 ± 15.1	0.287 ^a
Male		11.0 ± 13.5	12.2 ± 15.5	8.2 ± 6.0	0.896 ^a

^a Mann Whitney U test ^b Chi-square test ^c Fisher Exact test

MARS: Medical Adherence Report Scale-5

IES-R: Impact of Event Scale – Revised