

WEB TABLE 1 Characteristics of Children with Nephrotic Syndrome having COVID-19 Infection

Characteristics	Value
Age (y) ^a	6.20 (3.81-9.0)
Male sex	16 (53.3)
History of contact	14 (46.7)
Symptoms of COVID-19	
Fever	10 (33.3)
Cough	9 (30.0)
Fast breathing	4 (13.3)
Myalgia	4 (13.3)
Headache	3 (10.0)
Nausea and Vomiting	3 (10.3)
Abdominal Pain	7 (23.3)
Diarrhea	4 (13.3)
Severity of COVID-19	
mild/moderate/severe	12 (40)/11 (36.6)/(23.3)
Respiratory support	
Oxygen	8 (26.6)
Invasive Support	3 (10.0)
Outcome	
Discharge	25 (83.3)
Death	5 (16.6)
Type of Nephrotic Syndrome	
FENS/IFRNS/FRNS/SDNS/SRNS	7 (23.3)/ 3(10.0)/ 2(6.66)/ 8(26.6)/ 10(33.3)
DTTNS	12 (40)
Status of Nephrotic syndrome	
Relapse	23 (76.6%)
Remission	07 (30.4%)
Immunosuppression	
Steroids	29 (96.6)
Mycophenolate mofetil	07 (23.3)
Calcineurin inhibitor	07 (23.3)
Rituximab	02 (6.66)
Cyclophosphamide	03 (10.0)
Levamisole	04 (13.3)
Hospital stay (d) ^a	7.23 (7.5)
Baseline serum creatinine (mg/dL) ^a	0.37 (0.113)
Peak serum creatinine (mg/dL) ^a	0.50 (0.45)
AKI	9 (30)
Stage 1	1 (3.33)
Stage 2	4 (13.3)
Stage 3	4 (13.3)
Complications	
Pneumonia	9 (30.0)
Sepsis	17(56.6)
Shock and vasopressor support	3 (10.0)
Antibiotics	17 (56.6)
Nephrotoxic medications	9 (30.0)
Laboratory Results	
Hemoglobin (gm%) ^b	11.3 (2.57)
Total leukocyte count (mm ³) ^b	11000 (4283.24)
Absolute neutrophilic count (mm ³) ^b	5951 (8098.90)
PMN/Lymphocyte Ratio ^b	2.32 (2.12)
Platelet Count (mm ³) ^b	445000 (231387.7)

Values are expressed as n(%), or ^a (median IQR), or ^b (mean SD)

FENS, first episode nephrotic syndrome; IFRNS, Infrequent relapsing nephrotic syndrome; FRNS, frequent relapsing nephrotic syndrome; SDNS, steroid dependent nephrotic syndrome; SRNS, steroid resistant nephrotic syndrome; DTTNS, difficult to treat nephrotic syndrome; AKI, acute kidney injury; PMN, polymorph nuclear cell