

Table S4

Demographic and clinical characteristics of subgroups

		Backswitch group (N=28)	Inverse group (N=37)
Sex	Male	12 (42.9%)	14 (37.8%)
	Female	16 (57.1%)	23 (62.2%)
Age [in years]^a	Mean	77.6±1.4	79.5±1.6
	Median	76.9 (62.1;93.2)	79.7 (63.5;98.1)
Study eye	Right Eye	13 (46.4%)	16 (43.2%)
	Left Eye	15 (53.6%)	21 (56.8%)
Visual aquity^a [in LogMAR]	Mean	0.47±0.06	0.61±0.07
	Median	0.4 (0;1.5)	0.5 (0.1±1.5)
Intraocular pressure^a [in mmHg]	Mean	13.7±0.5	13.9±0.4
	Median	14 (9;20)	13.5 (9;18)
CRT [in μm]^a	Mean	329.4±18.7	290.1±12.1
	Median	301 (204;619)	290 (129;464)
Macular volume^a [in mm³]	Mean	8.2±0.24	7.9±0.18
	Median	8.17 (5.6;11.4)	7.9 (5.2;9.3)
Treatment time [in years]	Mean	5.5±0.5	4.2±0.4
	Median	5.9 (1.2;9.8)	4.1 (0.6;8.0)
IVT injections	Mean	34.6±2.6	25.5±2.1
	Median	33.5 (10;58)	24 (6;64)

^aBaseline = date of the last IVT injection before switch to bevacizumab

