

**Supplementary Table S1.** Number of subjects using antihypertensive agents and other medications.

Antihypertensive medications	Number of subjects
Calcium channel blockers	83
Beta blockers	75
Angiotensin II receptor blockers	68
Thiazide diuretics	60
Angiotensin converting enzyme inhibitors	45
Potassium sparing diuretics	21
Prazocin	17
Moxonidine	9
Furosemide	7
Beta + alpha blockers	3
Minoxidil	1
Other medications	Number of subjects
Vitamin D	64
Low acetylsalicylic acid	49
Antidepressant	36
Female hormones	35
Metformin	23
Proton pump inhibitor	22
Calcium supplement	22
Thyroxin*	22
Potassium supplement	18
Inhaled corticosteroids	17
Antihistamine	13
Tamsulosin or alfuzosin	13
Antirheumatic drugs	11
Insulin	10
Inhaled long-acting beta-agonist	9
Magnesium supplement	9
Oral anticoagulants	8
Benzodiazepine	7
Dipeptidyl peptidase-4 inhibitor	7
Allopurinol	7
Gabapentin or pregabalin	6
5-alpha reductase inhibitor	6
Intrauterine device	5
Nonsteroidal anti-inflammatory drug	5
Glaucoma medication	4

Ezetimibe	3
Glucagon-like peptide-1 agonist	3
Antiepileptic	3
Opioid	3
Sulphonyl urea	2
Triiodothyronine	2
Quetiapine	2
Clopidogrel	2
Fibrate	1
Sodium-glucose co-transporter-2 inhibitor	1
Amiodarone	1
Oxybutynin	1
Letrozole	1
Hydroxycarbamide	1
Dipyridamole	1
Aripiprazole	1
Levomepromazine	1
Pramipexol	1
Prednisolone	1

---

\*All subjects were euthyroid