Supplementary Table-1: Data tables of Figures 2 to 4

S1a. Figure-2 Data Table: % of patients with Intraoperative complications in cataract surgery in ten study hospitals (H1-H10)

Year	H1	H2	НЗ	H4	Н5	Н6	H7	Н8	Н9	H10	AVERAGE	SD	Coefficient of Variation
2012	1.71%	0.67%	0.79%	1.84%	0.77%	1.12%	1.74%	0.25%	1.66%	1.02%	1.2%	0.55%	48%
2013	1.36%	0.74%	0.62%	1.70%	0.98%	0.34%	2.08%	0.96%	0.24%	0.69%	1.0%	0.59%	60%
2014	1.10%	0.62%	0.49%	1.32%	0.93%	0.26%	2.61%	0.85%	0.34%	0.55%	0.9%	0.69%	76%
2015	1.19%	0.63%	0.56%	0.85%	0.72%	0.23%	1.98%	0.57%	0.37%	1.08%	0.8%	0.50%	62%
2016	1.27%	1.20%	0.54%	0.47%	0.53%	0.20%	1.43%	1.02%	0.46%	0.67%	0.8%	0.42%	54%
2017	1.46%	0.74%	0.56%	0.74%	0.46%	0.18%	1.14%	0.18%	0.63%	0.65%	0.7%	0.39%	58%
2018	1.28%	0.31%	0.58%	0.56%	0.51%	0.11%	0.72%	0.22%	0.17%	0.10%	0.5%	0.36%	80%
2019	1.51%	0.38%	0.59%	0.39%	0.58%	0.14%	0.54%	0.27%	0.29%	0.20%	0.5%	0.39%	80%
2020	1.39%	0.50%	0.42%	0.29%	0.52%	0.46%	1.40%	0.24%	0.19%	0.56%	0.6%	0.44%	73%

S1b. Figure-3 Data Table: % of patients with uncorrected visual acuity (>=6/12 [20/40]) at postoperative follow-up visit (2 to 7 weeks) in ten study hospitals (H1-H10)

Year	H1	H2	Н3	H4	Н5	Н6	Н7	Н8	Н9	H10	AVERAGE	SD	Coefficient of Variation
2012	84.0%	75.7%	72.2%	79.1%	85.0%	85.2%	86.5%	89.5%	76.5%	74.8%	80.8%	5.9%	7.3%
2013	86.4%	81.7%	77.8%	82.2%	82.8%	88.2%	88.2%	83.1%	82.3%	76.5%	83.5%	3.9%	4.7%
2014	87.9%	90.8%	84.2%	85.3%	86.6%	94.7%	88.9%	91.6%	84.9%	77.1%	87.2%	4.9%	5.6%
2015	86.7%	87.8%	87.2%	89.4%	89.1%	92.5%	88.9%	86.9%	87.2%	86.0%	87.9%	1.9%	2.1%
2016	90.8%	89.9%	88.8%	90.2%	90.7%	89.1%	90.1%	92.7%	89.4%	90.0%	90.2%	1.1%	1.2%
2017	91.1%	90.3%	86.3%	90.5%	88.2%	86.3%	87.4%	90.1%	88.4%	89.7%	89.4%	1.8%	2.0%
2018	89.8%	88.0%	86.8%	88.6%	88.7%	86.1%	86.3%	86.3%	90.7%	93.0%	88.4%	2.2%	2.5%
2019	84.2%	87.4%	86.9%	88.7%	87.6%	87.8%	84.5%	86.3%	89.1%	93.3%	87.6%	2.6%	2.9%
2020	85.3%	83.9%	88.6%	90.2%	92.0%	92.6%	89.2%	88.2%	92.2%	95.7%	89.8%	3.5%	4.0%

S1c. Figure-4 Data Table: % of patients with Spherical Equivalent (within ± 0.5D) at post-operative follow-up visit in ten study hospitals (H1-H10)

Year	H1	H2	Н3	H4	Н5	Н6	H7	Н8	Н9	H10	AVERAGE	SD	Coefficient of
													Variation
2012	86.3%	75.6%	71.9%	77.7%	80.6%	70.9%	84.7%	80.8%	73.9%	60.1%	76.3%	7.2%	9.5%
2013	85.6%	75.2%	74.7%	79.7%	80.9%	77.5%	88.1%	72.3%	67.7%	63.8%	76.6%	7.2%	9.5%
2014	87.7%	85.2%	80.3%	86.1%	84.4%	83.3%	88.0%	84.0%	73.6%	64.3%	81.7%	7.5%	9.2%
2015	90.9%	87.2%	84.6%	91.5%	90.4%	83.3%	87.1%	82.8%	76.3%	79.9%	85.4%	4.9%	5.7%
2016	91.9%	88.6%	84.9%	93.8%	92.7%	83.3%	88.0%	86.5%	77.6%	89.4%	87.7%	4.9%	5.6%
2017	91.4%	88.1%	84.0%	93.2%	91.3%	79.4%	88.6%	86.7%	80.8%	92.0%	87.6%	4.9%	5.6%
2018	90.1%	80.0%	86.2%	91.3%	90.9%	83.4%	86.9%	85.1%	88.6%	91.3%	87.4%	3.9%	4.5%
2019	87.8%	82.0%	86.5%	92.5%	88.9%	86.1%	81.1%	81.8%	84.7%	85.6%	85.7%	3.7%	4.3%
2020	85.5%	79.3%	85.3%	93.6%	92.9%	88.1%	84.0%	87.4%	84.7%	92.5%	87.3%	4.8%	5.5%