### Supplementary table 1

#### Main inclusion criteria

- 1. Underwent a radical hysterectomy in a European Institution for stage IB1 cervical cancer (FIGO 2009), from January 1, 2013 to December 31, 2014.
- 2. Patients age 18 years' old or older and
- 3. Histology must be either squamous cell carcinoma, adenosquamous carcinoma, or adenocarcinoma.
- 4. Patient must have received a preoperative pelvic MRI confirming
  - a. a tumor diameter less or equal than 4 cm
  - b. with no parametrial invasion and
- 5. Patient must have received a preoperative CT scan, MRI, or PET-CT without extracervical metastatic disease.
- 6. The operative report had to describe either a Type B–C radical hysterectomy with a bilateral pelvic lymphadenectomy, including at least a total of 10 pelvic nodes.
- 7. There had to be documentation of tumor size, margins, and nodal status.

#### Main exclusion cirteria

- 1. Any other histological type was diagnosed;
- 2. Tumor size larger than 4cm;
- 3. Evidence of past history of any invasive tumor, previous chemotherapy or radiation, suspicious lymph nodes, or metastatic disease on preoperative imaging;
- 4. Conversion from minimally invasive surgery to laparotomy;
- 5. Missing data on follow-up or adjuvant therapy;
- 6. Less than 10 pelvic lymph node count.
- 7. Missing data on performance of SNB+lymphadenectomy or lymphadenectomy

# Supplementary table 2

## Evaluation of compliance of Sedlis criteria

LVSI	Stromal invasion (deep 1/3)	Tumor diameter <u>&gt;</u> 2 cm	SEDLIS CRITERIA
+	deep	-	Yes
+	middle	+	Yes