

Determination of lindane in surface water samples and its degradation by hydrogen peroxide and persulfate assisted TiO₂-based photocatalysis

Sanaullah Khan¹, Javed Ali Khan^{2,*}, Noor S. Shah³, Murtaza Sayed⁴, Muhammad Ateeq²,
Sabah Ansar⁵, Grzegorz Boczkaj^{6,7}, Umar Farooq^{8,9}

¹Departmen of Chemistry, Women University Swabi, 23430, Pakistan

²Department of Chemistry, Abdul Wali Khan University Mardan, Mardan 23200, Pakistan

³Department of Environmental Sciences, COMSATS University Islamabad, Vehari Campus,
61100, Pakistan

⁴Radiation Chemistry Laboratory, National Centre of Excellence in Physical Chemistry,
University of Peshawar, Peshawar 25120, Pakistan

⁵Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, King Saud
University, P.O. Box 10219, Riyadh, 11433, Saudi Arabia

⁶Department of Sanitary Engineering, Faculty of Civil and Environmental Engineering,
Gdansk University of Technology, G. Narutowicza St. 11/12, 80-233, Gdansk, Poland

⁷EkoTech Center, Gdansk University of Technology, G. Narutowicza St. 11/12, 80-233,
Gdansk, Poland

⁸Department of Chemistry, COMSATS University Islamabad, Abbottabad-Campus 22060,
Abbottabad, Pakistan

⁹Beijing National Laboratory for Molecular Sciences, State Key Laboratory of Molecular
Reaction Dynamics, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190,
China

*Corresponding author e-mails: javedkhan@awkum.edu.pk; khanjaved2381@gmail.com
(J.A. Khan)

Fax: +92-937-542189; Tel: +92-937-929122

Table S1. Physico-chemical parameters of the water sample collected from Amangarh, Nowshera (N7) where maximum concentration of lindane ($2.60 \mu\text{g L}^{-1}$) was detected.

pH	Conductivity (μS)	TOC (mg/L)	TH (mg/L)	T. alkalinity (mg/L)	TSS (mg/L)	TDS (mg/L)	Nitrates (mg/L)	Chlorides (mg/L)	Sulphates (mg/L)	Turbidity (NTU)
7.3	712.8	4.58	286.5	172.5	245.6	368.4	3.4	356.3	86.5	17.6

*T (Total); TOC (Total organic carbon); TH (Total Hardness); TTS (Total suspended solids); TDS (Total dissolved solids)

Table S2. Method validation of SPME followed by GC-ECD, in terms of R²-value, LOD and LOQ.

Analyte	Equation	Correlation coefficient (R ²)	LOD ($\mu\text{g L}^{-1}$)	LOQ ($\mu\text{g L}^{-1}$)
Lindane	$y = 66.71 x + 2.27$	0.997	0.008	0.037

Table S3. Cultivation area, cotton production, yield value and pesticides consumed in Pakistan during 1997-2006 (Economic Survey of Pakistan; 2003-2004 and 2005-2006).

Year	Area (Thousand hectares)	Production (Thousand bales)	Yield (kg/ha)	Total pesticides in tones (Imported + Locally manufactured)
1997-98	2960	9184	528	42,732.0
1998-99	2923	8790	512	44,872.0
1999-00	2983	11,240	641	45,232.0
2000-01	2927	10,732	570	44,762.0
2001-02	3116	10,613	579	46,132.0
2002-03	2794	10,211	622	46,943.0
2003-04	2989	10,048	572	47,432.0
2004-05	3193	14,265	760	46,572.0
2005-06	3096	12,417	682	46,951.0

Table S4. Effect of different scavengers on k_{obs} of lindane by solar/TiO₂. Experimental conditions: [Lindane]₀ = 2.60 µg L⁻¹; [MeOH]₀ = [*i*-PrOH]₀ = [BQ]₀ = 50 mM; [PDC]₀ = 100 µM; Mass of TiO₂ film = 9.02 mg; Thickness of TiO₂ film = 1.02 µm; Area of TiO₂ film = 3750 mm²; pH = 7.3.

Scavenger	$k_{\text{obs}} (\text{h}^{-1})$
Control (No scavenger)	0.0309
Iso-PrOH (<i>i</i> -PrOH)	0.0126
Methanol (MeOH)	0.0089
Benzoquinone (BQ)	0.0258
Potassium dichromate (PDC)	0.0230

Table S5. Acute and chronic toxicities of lindane and its detected DPs towards fish, daphnia

Compound	Acute toxicity (mg/L)			Chronic toxicity (mg/L)		
	LC ₅₀	LC ₅₀	EC ₅₀	ChV	ChV	ChV
	Fish	Daphnia	Green Algae	Fish	Daphnia	Green Algae
Lindane	2.24	1.57	2.76	0.280	0.273	1.15
HCB	0.08	0.065	0.211	0.012	0.017	0.122
TCB	2.77	1.88	2.93	0.334	0.301	1.14
DCB	8.52	5.45	6.62	0.958	0.738	2.26
DCP	25.5	15.6	15.8	2.72	1.87	4.87
CBD	232.0	128.0	84.0	21.9	11.5	20.6

and green algae.