

Supplementary table S4. Functional class by paediatric and WHO systems at the time of diagnosis of pulmonary hypertension, by WSPH group. Column percentages for each system total to 100%. The p-value tests equality of the distributions of each system for the Group 1 vs. Group 3 cohorts.

Functional Class	Overall (n=1475)	WSPH					Grp 1 vs. 3 Pvalue
		Group 1 (n=663)	Group 2 (n=54)	Group 3 (n=720)	Group 4 (n=8)	Group 5 (n=30)	
Pediatric	791	341	28	400	791	17	<.001
I	104 (13.1%)	55 (16.1%)	7 (25.0%)	39 (9.8%)	0 (0%)	3 (17.6%)	
II	229 (29.0%)	122 (35.8%)	7 (25.0%)	90 (22.5%)	2 (40.0%)	8 (47.1%)	
III A	154 (19.5%)	81 (23.8%)	9 (32.1%)	62 (15.5%)	0 (0%)	2 (11.8%)	
III B	65 (8.2%)	28 (8.2%)	2 (7.1%)	33 (8.3%)	1 (20.0%)	1 (5.9%)	
IV	239 (30.2%)	55 (16.1%)	3 (10.7%)	176 (44.0%)	2 (40.0%)	3 (17.6%)	
World Health Organization	503	293	25	172	503	10	0.011
I	105 (20.9%)	49 (16.7%)	4 (16.0%)	48 (27.9%)	0 (0%)	4 (40.0%)	
II	204 (40.6%)	132 (45.1%)	12 (48.0%)	55 (32.0%)	2 (66.7%)	3 (30.0%)	
III	138 (27.4%)	80 (27.3%)	7 (28.0%)	49 (28.5%)	0 (0%)	2 (20.0%)	
IV	56 (11.1%)	32 (10.9%)	2 (8.0%)	20 (11.6%)	1 (33.3%)	1 (10.0%)	