

Supplementary Material

Use of Intranasal Rescue Therapy for Seizure Clusters in Students With Epilepsy: Nurses' Knowledge, Perceptions, and Practice

Authors: Nancy Santilli, PNP, MN^a; Sandra Dewar, APRN, PhD^b; Cynthia Guerra, MBA^c;

Sunita N. Misra, MD, PhD^c; Adrian L. Rabinowicz, MD^c; Enrique Carrazana, MD^{c,d}

Supplementary Table 1. Who Did You or Will You Train on the Use of Intranasal Diazepam?

Answer	Response Rate, n (%)
	n=88
School nurses	55 (62.5)
Teachers	54 (61.4)
School aides	50 (56.8)
Administration staff	46 (52.3)
Coaches	27 (30.7)
Other	17 (19.3)

Supplementary Table 2. Of All the Times a Student Required Intranasal Diazepam, How Often Did You Administer It to the Student?

Answer	Response Rate, n (%)^a
	n=43
All of the time	28 (65.1)
75% of the time	0 (0)
50% of the time	4 (9.3)
25% of the time	3 (7.0)
≤24% of the time	8 (18.6)

^a“N/A” responses were excluded from tabulation.

Supplementary Table 3. If the Student Would Be Allowed to Participate in Off School Premises or Extracurricular Activities, Who Would Be Required to Carry Intranasal Diazepam for the Student?

Answer	Response Rate, n (%)
	n=190
Nurse	40 (21.1)
Parent/guardian	21 (11.1)
Teacher/coach	90 (47.4)
Other	39 (20.5)