

Supplemental Material for

Generalization of contextual fear is sex-specifically affected by high salt intake

Jasmin N. Beaver^{1,2}, Brady L. Weber^{1,2}, Matthew T. Ford¹, Anna E. Anello^{1,2}, Kaden M. Ruffin¹, Sarah K. Kassis^{1,2}, T. Lee Gilman^{1,2,3*}

¹Department of Psychological Sciences, Kent State University, Kent, Ohio, United States of America

²Brain Health Research Institute, Kent State University, Kent, Ohio, United States of America

³Healthy Communities Research Institute, Kent State University, Kent, Ohio, United States of America

*Corresponding Author

Email: lgilman1@kent.edu (TLG)

S13. Three-way ANOVAs on log-transformed serum corticosterone levels in control no shock mice across Experiments.

Corticosterone	No Shock Groups Across Experiments		
Sex	F(1,81)=12.77	p<0.001	partial η^2 =0.136
Diet	F(1,81)=2.498	p=0.118	partial η^2 =0.030
Experiment	F(2,81)=12.81	p<0.001	partial η^2 =0.240
Sex × Diet	F(1,81)=1.654	p=0.202	partial η^2 =0.020
Sex × Experiment	F(2,81)=3.498	p=0.035	partial η^2 = 0.080
Diet × Experiment	F(2,81)=0.527	p=0.592	partial η^2 =0.013
Sex × Diet × Experiment	F(2,81)=1.512	p=0.227	partial η^2 =0.036