

Supplementary Material

Nicotinamide mononucleotide (NMN) alleviates the poly(I:C)-induced inflammatory response in human primary cell cultures

Hitomi Sano, Anton Kratz, Taiko Nishino, Haruna Imamura, Yuki Yoshida, Noriaki Shimizu, Hiroaki Kitano and Ayako Yachie

1 *List of Supplementary Figures*

Supplementary Figure 1. Images of representative cells from each condition at the time of RNA isolation.

Supplementary Figure 2. PCA plot for FPKM transcriptome data with poly(I:C) (2.5 µg/mL) induction (red) and without poly(I:C) induction (blue) in HPMEC (A) and HCAEC (B).

Supplementary Figure 3. Sankey diagram illustrating common genes between HPMEC and HCAEC.

Supplementary Figure 4. Boxplots for additional eigengenes and list of pathways overrepresented in selected modules in HPMEC (A) and HCAEC (B).

Supplementary Figure 5. Gene expression of the seven sirtuin genes in HPMEC cells under different poly(I:C) conditions and NMN doses.

Supplementary Figure 6. Gene expression of the seven sirtuin genes in HCAEC cells under different poly(I:C) conditions and NMN doses.

2 *List of Supplementary Tables*

Supplementary Table 1. List of primers.

Supplementary Table 2. PCA loadings.

Supplementary Table 3. Traits for WGCNA.

3 *List of Supplementary Information*

Supplementary Information 1.

List of top 20 genes up-regulated by high-NMN in poly(I:C)-induced HPMEC.

List of top 20 genes down-regulated by high-NMN in poly(I:C)-induced HPMEC.

List of top 20 genes down-regulated by high-NMN in poly(I:C)-induced HCAEC.

List of top 20 genes up-regulated by high-NMN in poly(I:C)-induced HCAEC.

Supplementary Information 2.

Gene Ontology pathways enriched in up- and downregulated genes of HPMEC and HCAEC.

Supplementary Information 3.

Module-trait relationships (WGCNA p-values) in HPMEC.

Module-trait relationships (WGCNA p-values) in HCAEC.

Gene Ontology pathways overrepresented in the p-sienna3, p-green, c-magenta, c-purple modules. These are the full lists of which only excerpts are shown in Figure 2 E, F.

Supplementary Information 4.


Gene Ontology pathways overrepresented in the p-purple, c-yellow modules. These are the full lists of which only excerpts are shown in Figure 3 C, D.

Supplementary Information 5.


Gene Ontology pathways overrepresented in the p-lightgreen, c-brown modules. These are the full lists of which only excerpts are shown in Supplementary Figure 4.

Supplementary Information 6.


Gene Ontology pathways overrepresented in the p-darkred, c-black modules. These are the full lists of which only excerpts are shown in Figure 5 A, B.





Supplementary Figure 1: Images of representative cells from each condition at the time of RNA isolation. Yellow bar 100 μ m.










Supplementary Figure 2: PCA plot for FPKM transcriptome data with poly(I:C) (2.5 μ g/mL) induction (red) and without poly(I:C) induction (blue) in HPMEC (A) and HCAEC (B). Pp: Poly(I:C)+. Pm: Poly(I:C)-. low: 0.1 mM, mid: 1.0 mM, high: 10 mM NMN.










Supplementary Figure 3: Sankey diagram illustrating common genes between HPMEC and HCAEC.



Supplementary Figure 4: Boxplots for eigengenes and list of pathways overrepresented in selected modules in HPMEC (A) and HCAEC (B). Pp: Poly(I:C)+. Pm: Poly(I:C)-. low: 0.1 mM, medium: 1.0 mM, high: 10 mM NMN.



Supplementary Figure 5: Gene expression of the seven sirtuin genes in HPMEC cells under different poly(I:C) conditions and NMN doses. Pp: Poly(I:C)+. Pm: Poly(I:C)-. low: 0.1 mM, med: 1.0 mM, high: 10 mM NMN.



Supplementary Figure 6: Gene expression of the seven sirtuin genes in HCAEC cells under different poly(I:C) conditions and NMN doses. Pp: Poly(I:C)+. Pm: Poly(I:C)-. low: 0.1 mM, med: 1.0 mM, high: 10 mM NMN.