

# **SUPPLEMENTAL MATERIAL**

**Table S1. Patient Characteristics by Race and Ethnicity, and Obstructive versus Nonobstructive Phenotype**

	Obstructive HCM			p	Nonobstructive HCM			p
	White	Black	Hispanic		White	Black	Hispanic	
<b>Total (N)</b>	18494 (77.0%)	4106 (17.1%)	1410 (5.80%)		20331 (69.8%)	6763 (23.2%)	2013 (6.90%)	
<b>Female</b>	11216 (60.6%)	2737 (66.7%)	854 (60.6%)	<0.001	11203 (55.1%)	3561 (52.7%)	1063 (52.8%)	<0.001
<b>Age</b>				<0.001				<0.001
<b>18-34</b>	530 (2.90%)	260 (6.30%)	102 (7.20%)		672 (3.30%)	460 (6.80%)	154 (7.70%)	
<b>35-54</b>	2974 (16.1%)	1073 (26.1%)	394 (27.9%)		3432 (16.9%)	2058 (30.4%)	687 (34.1%)	
<b>55-64</b>	3488 (18.9%)	977 (23.8%)	278 (19.7%)		3513 (17.3%)	1543 (22.8%)	337 (16.7%)	
<b>65-74</b>	4647 (25.1%)	896 (21.8%)	279 (19.8%)		4776 (23.5%)	1330 (19.7%)	356 (17.7%)	
<b>75+</b>	6855 (37.1%)	900 (21.9%)	357 (25.3%)		7938 (39.0%)	1372 (20.3%)	479 (23.8%)	
<b>Insurance</b>				<0.001				<0.001
<b>Medicare</b>	12286 (66.4%)	2290 (55.8%)	713 (50.6%)		13667 (67.2%)	3626 (53.6%)	945 (46.9%)	
<b>Medicaid</b>	1258 (6.8%)	820 (20.0%)	308 (21.8%)		1507 (7.40%)	1454 (21.5%)	569 (28.3%)	
<b>Private insurance</b>	4622 (25.0%)	813 (19.8%)	318 (22.6%)		4777 (23.5%)	1348 (19.9%)	396 (19.7%)	
<b>Uninsured</b>	328 (1.80%)	183 (4.50%)	71 (5.00%)		380 (1.90%)	335 (5.00%)	103 (5.10%)	
<b>Median ZIP income</b>				<0.001				<0.001
<b>Q1: \$0-\$47,999</b>	3927 (21.2%)	2093 (51.0%)	474 (33.6%)		5625 (27.7%)	792 (11.7%)	357 (17.7%)	
<b>Q2: \$48,000-\$60,999</b>	4744 (25.7%)	891 (21.7%)	351 (24.9%)		5423 (26.7%)	1169 (17.3%)	425 (21.1%)	
<b>Q3: \$61,000-\$81,999</b>	4791 (25.9%)	662 (16.1%)	351 (24.9%)		5140 (25.3%)	1472 (21.8%)	499 (24.8%)	
<b>Q4: \$82,000+</b>	5032 (27.2%)	460 (11.2%)	234 (16.6%)		4143 (20.4%)	3330 (49.2%)	732 (36.4%)	
<b>Rurality</b>				<0.001				<0.001
<b>Urban</b>	9407 (50.9%)	2927 (71.3%)	1074 (76.2%)		10213 (50.2%)	4579 (67.7%)	1355 (67.3%)	
<b>Town</b>	5815 (31.4%)	920 (22.4%)	268 (19.0%)		6534 (32.1%)	1710 (25.3%)	530 (26.3%)	
<b>Rural</b>	3272 (17.7%)	259 (6.30%)	68 (4.80%)		3584 (17.6%)	474 (7.00%)	128 (6.40%)	
<b>Comorbidities</b>								
<b>Hypertension, complicated</b>	5340 (28.9%)	1832 (44.6%)	433 (30.7%)	<0.001	6556 (32.2%)	3489 (51.6%)	669 (33.2%)	<0.001
<b>Cardiac arrhythmia</b>	11507 (62.2%)	2097 (51.1%)	807 (57.2%)	<0.001	12496 (61.5%)	3098 (45.8%)	988 (49.1%)	<0.001
<b>Congestive heart failure</b>	7805 (42.2%)	1913 (46.6%)	567 (40.2%)	<0.001	9099 (44.8%)	3406 (50.4%)	837 (41.6%)	<0.001
<b>Obesity</b>	3479 (18.8%)	1008 (24.5%)	302 (21.4%)		3514 (17.3%)	1528 (22.6%)	394 (19.6%)	<0.001

<b>Diabetes</b>	1728 (9.30%)	683 (16.6%)	176 (12.5%)		2166 (10.7%)	1267 (18.7%)	328 (16.3%)	<0.001
<b>Presence of ICD</b>	2321 (12.6%)	630 (15.3%)	209 (14.8%)	<0.001	2657 (13.1%)	755 (11.2%)	202 (10.0%)	<0.001

HCM, hypertrophic cardiomyopathy; ICD, implantable cardioverter-defibrillator. Comparisons were conducted using chi-square tests of independence.

**Table S2. Hospital-Year Characteristics**

	<b>Hospital-Years</b>
<b>Total (N)</b>	13020
<b>Size</b>	
<b>Small</b>	3429 (26.3%)
<b>Medium</b>	4081 (31.3%)
<b>Large</b>	5510 (42.3%)
<b>Ownership</b>	
<b>Government, nonfederal (public)</b>	1290 (9.9%)
<b>Private, not-for-profit (voluntary)</b>	9726 (74.7%)
<b>Private, investor owned (proprietary)</b>	2004 (15.4%)
<b>Teaching status</b>	
<b>Rural</b>	1734 (13.3%)
<b>Urban - Non-teaching hospital</b>	4498 (34.5%)
<b>Urban - Teaching hospital</b>	6788 (52.1%)
<b>Region</b>	
<b>Northeast</b>	2562 (19.7%)
<b>Midwest</b>	3137 (24.1%)
<b>South</b>	4715 (36.2%)
<b>West</b>	2606 (20.0%)

Because hospitals cannot be tracked year to year, data are presented as hospital-years. To estimate annual n, divide by 7.

**Table S3. In-hospital Mortality Rate by Hypertrophic Cardiomyopathy Procedure**

	<b>Mortality Rate</b>
<b>Overall (n=53117)</b>	3.23%
<b>Septal myectomy (n=1654)</b>	3.33%
<b>Alcohol septal ablation (n=772)</b>	1.55%
<b>ICD procedure (n=1598)</b>	0.56%
<b>Any HCM procedure (n=3871)</b>	1.91%

Figure S1. Participant Flow Diagram

