

Birth

Please complete the survey below.

Thank you!

-
- 1) Timing of Delivery:
The decision for timing of birth should be made by :
- Obstetrics
 CIF team
 MDT including obstetric and CIF team
 I don't know
-
- 2) At what gestation should delivery be considered in the absence of fetal compromise?
- < 37 weeks
 37-40weeks
 >40 weeks
 I don't know
-
- 3) The decision to define the mode of infant delivery should be individualised according to patient specific factors?
- Strongly agree
 Agree
 Neutral
 Disagree
 Strongly disagree
 I don't know

Please specify patient factors that would mean you would recommend caesarean section, rather than vaginal delivery:

	Strongly agree	Agree	Neutral	Disagree	Strongly disagree	I don't know
4) Absence of a stoma	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
5) Presence of perianal disease	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
6) Multiple abdominal surgeries	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
7) Obstetric indications only	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
8) CIPO diagnosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
9) Presence of ileo-anal pouch	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
10) Poor abdominal wall strength	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>