

Opinions

Please complete the survey below.

Thank you!

In your opinion or experience, is the fertility of women receiving home parenteral support generally:

- Increased
- Decreased
- Same as general population
- My experience is too small to make reliable comment
- It depends on the patient's underlying disease or clinical condition'

In your opinion or experience, is the fertility of men receiving home parenteral support

- Increased
- Decreased
- Same as general population
- My experience is too small to make reliable comment
- It depends on the patient's underlying disease or clinical condition

In your opinion should women with CIF be advised against pregnancy?

- Yes
- No
- No opinion

Unless brought up by your patient, would you routinely discuss contraception for Home Parenteral Nutrition-dependent patients?

- Yes
- No

If Yes. Which option best represents your opinion or standard recommendation on contraception for HPN-dependent patients?

- No contraception or left to patient choice
- Routinely recommend a single method
- Recommend two forms of contraception e.g, hormonal and barrier

In your opinion or experience, are fetal abnormalities for women receiving home parenteral support:

- More frequent
- Less frequent
- Same as general population
- My experience is too small to make reliable comment

In your opinion or experience, are pregnancy-related maternal complications, in women receiving treatment with home parenteral support:

- More frequent
- Less frequent
- Same as general population
- My experience is too small to make reliable comment

In your opinion or experience, preterm delivery for women receiving home parenteral support is:

- More frequent
- Less frequent
- Same as general population
- My experience is too small to make reliable comment

In your opinion or experience, the infant birth weight corrected for gestation at delivery seen for women receiving home parenteral support is:

- Above average
- Below Average
- Same as general population
- My experience is too small to make reliable comment

HPN specific complications.**In your opinion or experience is a pregnant women more likely to have a CIF related complication?**

	Strongly agree	Agree	Neutral	Disagree	Strongly disagree
Catheter related blood stream infection (CRBSI)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
CVC associated thrombosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Intestinal failure associated liver disease (IFALD)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Please rank the underlying intestinal failure disease according to its impact on the risk to conception. (1- highest risk 8- lowest risk)

	1	2	3	4	5	6	7	8
CIPO	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Crohn's disease	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Mesenteric ischaemia	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Mucosal disease e.g. coeliac	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Radiation enteritis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Scleroderma	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Surgical complication	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Immunodeficiency disorder	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Please rank the underlying intestinal failure disease according to its impact on the risk of a successful pregnancy. (1- highest risk and 8- lowest risk)

	1	2	3	4	5	6	7	8
CIPO	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Crohn's disease	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Mesenteric ischaemia	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Mucosal disease e.g. coeliac	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Radiation enteritis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Scleroderma	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Surgical complication	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Immunodeficiency disorders	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Do you routinely check Essential Fatty Acid (EFA) levels as part of pregnancy management in chronic intestinal failure? Yes No

If yes, at what points would you typically check them?

- Pre-conception
 Pregnancy
 Post-natal
 During duration of breastfeeding

In your opinion, would a specialist IF-pregnancy service improve care/outcomes for pregnant patients with chronic intestinal failure? Yes No No opinion