

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Extreme Temperature Exposure and Risks of Preterm Birth Subtypes Based on a Nationwide Survey in China

Guoqi Yu, Ling Yang, Ming Liu, Cuiping Wang, Xiaoli Shen, Lichun Fan, and Jun Zhang

Table of Contents

Table S1. The selection basis of the China Labor and Delivery Survey to determine the sample size of deliveries in each hospital according to their levels and types.

Table S2. Basic characteristics of climate and air pollution by outcomes in the China Labor and Delivery Survey (N=70,818).

Table S3. Basic characteristics of temperature exposure by delivery hospitals and geographic regions in the China Labor and Delivery Survey (N=70,818).

Table S4. Extreme climate event exposure by time periods and outcomes in the China Labor and Delivery Survey (N=70,818).

Table S5. Associations of extreme climate events with early and late preterm birth in the China Labor and Delivery Survey.

Table S6. Associations of extreme climate events with early and late preterm birth in the China Labor and Delivery Survey, stratified by clinical subtypes.

Table S7. Associations of extreme climate events with preterm birth in the China Labor and Delivery Survey, stratified by geographic regions.

Table S8. Associations of extreme climate events with preterm birth in the China Labor and Delivery Survey, stratified by child sex.

Table S9. Estimates from the Cox regression investigating associations of temperature exposure with PTB in the China Labor and Delivery Survey (N=70,818).

Table S10. Associations of extreme climate events with preterm birth under different temperature definitions in the China Labor and Delivery Survey.

Table S11. Associations of extreme climate events with subtypes of preterm birth during 3 and 4 weeks before delivery in the China Labor and Delivery Survey.

Table S12. Associations of extreme temperature with preterm birth risks during cold and warm seasons in the China Labor and Delivery Survey.

Table S13. Associations of extreme climate events with preterm birth among subjects without history of preterm birth in the China Labor and Delivery Survey, stratified by early and late preterm birth (N=69,970).

Table S14. Associations of extreme climate events with preterm birth among subjects without severe maternal disease in the China Labor and Delivery Survey, stratified by early and late preterm birth (N=69,883).

Table S15. Associations of extreme climate events with preterm birth in the China Labor and Delivery Survey, additionally adjusting for background temperature exposure (N=70,818).

Table S16. Associations of extreme climate events with preterm birth based on data with multiple imputation in the China Labor and Delivery Survey (N=70,818).

Figure S1. The geographical location of study areas in China.

Figure S2. Flowchart for population selection.

Figure S3. Directed acyclic graph for covariate selection.

Figure S4. Relationships between the average temperature exposed during different trimesters and the preterm birth risk in the China Labor and Delivery Survey (N=70,818) (See Tables S9 for corresponding numeric data).

References