

Table S1. Medications pre- and post-SGLT2 inhibitor use.

	Case 1	Case 2	Case 3	Case 4
Medications (Pre-SGLT2 inhibitor)	Amlodipine, aspirin, atorvastatin, carvedilol, ferrous sulfate, levothyroxine, losartan, Mag-tab SR, melatonin, mirtazapine, mycophenolate mofetil, sulfamethoxazole- trimethoprim, tacrolimus	Amlodipine, apixaban, fiber, levothyroxine, losartan, magnesium gluconate, sodium chloride, tolterodine tartrate, ergocalciferol, amiloride	Afatinib, sodium chloride, folic acid, midodrine, famotidine, amiloride, magnesium oxide, levothyroxine	Magnesium extended release, eplerenone, baclofen, calcium citrate/Vitamin D, Biotin, multivitamin, gabapentin, fexofenadine, albuterol inhaler
Medications (Post-SGLT2 inhibitor)	Amlodipine, aspirin, atorvastatin, carvedilol, ferrous sulfate, levothyroxine, losartan, melatonin, mirtazapine, mycophenolate mofetil, sulfamethoxazole- trimethoprim, tacrolimus	Amlodipine, apixaban, fiber, levothyroxine, losartan, magnesium gluconate, sodium chloride, tolterodine tartrate, ergocalciferol, amiloride	Afatinib, sodium chloride, folic acid, midodrine, famotidine, amiloride, magnesium oxide, levothyroxine, oxycodone, apixaban, fludrocortisone, sodium bicarbonate	Magnesium extended release, eplerenone, baclofen, calcium citrate/Vitamin D, Biotin, multivitamin, gabapentin, fexofenadine, albuterol inhaler

SGLT2, Sodium-glucose cotransporter-2