

Supporting Information




Fig. S1 (Linked to Fig.1). Histogram plots of fitness for all *L. pneumophila* genes represented on Tn-seq.

Histogram of WT (SK01) Tn-seq pool following either infection in BMDM (A) or growth in nutrient-rich AYE medium (B). Histogram of Δsde (SK02) Tn-seq pool following infection in BMDM (C) or growth in nutrient-rich AYE medium (D).




Fig. S2 (Linked to Fig.3). The integrity of LCVs harboring $\Delta sdhA$ strains after challenge with *L. pneumophila*.

Percent cytosolic bacteria was quantified based on antibody accessibility. BMDMs were infected with either WT or $\Delta sdhA$ strains for 2, 4, and 6 hr, fixed, and stained with antibodies. The internalized bacteria in the absence of permeabilization were calculated relative to the total infected population (mean \pm SEM; three biological replicates were performed and 100 LCVs were counted per biological replicate). Statistical analysis was conducted using unpaired two-tailed Student's t test (ns, not significant; ***p < 0.001).




Fig. S3 (Linked to Fig. 3). Transcription of *sde* genes is downregulated during *L.*

***pneumophila* infection of BMDMs.**

Transcript abundance of indicated genes was determined during infection. PMA-differentiated U937 cells were challenged with *L. pneumophila* WT and RNA was extracted at the noted time points. Transcripts were normalized to 16s rRNA, and then displayed relative to transcription level measured at 1hr post-infection and represented as fold change.




Fig. S4 (Linked to Fig. 3). The loss of *sdhA*, *ridL* and *legA3* aggravated vacuole disruption in Δsde strain.

Vacuole integrity was measured based on antibody accessibility. BMDMs in a 96 well plate were infected with the indicated strains for 2 hr, fixed and stained with antibodies. The images were taken using a Lionheart automated microscope with a 10X magnification objective. The internalized bacteria in the absence of permeabilization were calculated relative to total infected population to determine fraction of disrupted vacuoles (mean \pm SEM; three biological replicates were performed and 1000-3000 LCVs were counted per biological replicate). Statistical significance was tested using one-way ANOVA with Tukey's multiple comparisons; ***p <0.001.




Fig. S5 (Linked to Fig. 3). Uptake efficiency of *L. pneumophila* is independent of genetic background.

BMDMs were challenged with noted strains, washed 3X with PBS at 10 or 60 min post-infection, lysed and cell lysates were plated on CYE plates. The efficiency of association was determined with CFUs in the inoculum and after either 10 or 60 min. incubation with BMDMs (mean \pm SEM; two biological replicates were performed with three technical replicates per biological replicate). Statistical significance was determined using one-way ANOVA with Tukey's multiple comparisons; ns, non-significant.

Table S1. Strains, Plasmids and Oligonucleotides used in this study

Strains			
Name	Genotype	Description	Reference
<i>L.pneumophila</i>			
Lp02	Philadelphia 1 <i>thyA</i> ⁺ <i>rpsL</i> <i>hsdR</i>	Wild type strain	(1)
SK01	Lp02 <i>thyA</i> ⁺	Wild type strain <i>thyA</i> ⁺	This work
Lp03	<i>thyA</i> ⁺ <i>rpsL</i> <i>hsdR</i> <i>dotA03</i>	Icm/Dot translocation deficient	(1)
JV6113	Lp02 <i>ΔsidE</i> <i>ΔsdeC</i> <i>ΔsdeBA</i> (<i>Δlpg0234</i> <i>Δlpg2153</i> <i>Δlpg2156-2157</i>)	<i>sidE</i> family deletion mutant	(2)
SK02	JV6113 <i>thyA</i> ⁺	JV6113 strain <i>thyA</i> ⁺	This work
SK03	Lp02 <i>thyA</i> ⁺ <i>ΔsdhA</i>	<i>sdhA</i> deletion mutant	This work
SK04	JV6113 <i>thyA</i> ⁺ <i>ΔsdhA</i>	<i>sdhAsidE</i> family deletion mutant	This work
SK05	Lp02 <i>thyA</i> ⁺ <i>ΔridL</i>	<i>ridL</i> deletion mutant	This work
SK06	JV6113 <i>thyA</i> ⁺ <i>ΔridL</i>	<i>ridLsidE</i> family deletion mutant	This work
SK07	Lp02 <i>thyA</i> ⁺ <i>ΔlegA3</i>	<i>legA3</i> deletion mutant	This work
SK08	JV6113 <i>thyA</i> ⁺ <i>ΔlegA3</i>	<i>legA3sidE</i> family deletion mutant	This work
SK09	Lp02 <i>thyA</i> ⁺ <i>ΔsdhA</i> <i>ΔridL</i>	<i>sdhAridL</i> deletion mutant	This work
SK10	Lp02 <i>thyA</i> ⁺ <i>ΔsdhA</i> <i>ΔlegA3</i>	<i>sdhA legA3</i> deletion mutant	This work
SK11	Lp02 <i>thyA</i> ⁺ <i>ΔridL</i> <i>ΔlegA3</i>	<i>ridL legA3</i> deletion mutant	This work
SK12	Lp02 <i>thyA</i> ⁺ <i>kan</i> ^R <i>P_{ahpc}::lux</i>	wild type strain Lux ⁺	This work
SK13	JV6113 <i>thyA</i> ⁺ <i>kan</i> ^R <i>P_{ahpc}::lux</i>	<i>sidE</i> family deletion mutant Lux ⁺	This work
SK14	SK02 <i>kan</i> ^R <i>P_{ahpc}::lux</i>	<i>sdhA</i> deletion mutant Lux ⁺	This work
SK15	SK03 <i>kan</i> ^R <i>P_{ahpc}::lux</i>	<i>sdhAsidE</i> family deletion mutant Lux ⁺	This work
SK16	SK04 <i>kan</i> ^R <i>P_{ahpc}::lux</i>	<i>ridL</i> deletion mutant Lux ⁺	This work
SK17	SK05 <i>kan</i> ^R <i>P_{ahpc}::lux</i>	<i>ridLsidE</i> family deletion mutant	This work
SK18	SK06 <i>kan</i> ^R <i>P_{ahpc}::lux</i>	<i>legA3</i> deletion mutant Lux ⁺	This work
SK19	SK07 <i>kan</i> ^R <i>P_{ahpc}::lux</i>	<i>legA3 sidE</i> family deletion mutant Lux ⁺	This work
Lp03 lux ⁺	Lp03 <i>kan</i> ^R <i>P_{ahpc}::lux</i>	Icm/Dot translocation deficient Lux ⁺	(3)
JV4487	<i>ΔsidJ</i>	<i>sidJ</i> deletion mutant	(2)
SK20	Lp02 <i>ΔsdhA</i>	<i>sdhA</i> deletion mutant	This work
SK21	Lp02 <i>ΔsdhA</i> <i>ΔsidJ</i>	<i>sdhAsidJ</i> deletion mutant	This work
SK22	SK01+ pMMB207Δ267		This work
SK23	SK02+ pMMB207Δ267		This work
SK24	SK03+ pMMB207Δ267		This work
SK25	SK04+ pMMB207Δ267		This work

SK26	SK04+ pSK04	This work
SK27	SK04+ pSK05	This work
SK28	SK04+ pSK06	This work
SK29	SK04+ pSK07	This work
SK30	JV6113 $\Delta sdhA$	This work
SK31	SK29+ pSdeB _{WT}	This work
SK32	SK29+ pSdeB _{C118S}	This work
SK33	SK29+ pSdeB _{H416A}	This work
SK34	SK29+ pSdeB _{E859A}	This work

Plasmids

Name	Features	Description	Reference
pTO100MmeI	R6Kori <i>kan</i> ^R , <i>sacB</i> , <i>ampR</i> , <i>himar1</i> -MmeI, C9 transposase	Tn-seq transposon mutagenesis plasmid	(4)
pSR47S	R6Kori <i>sacB</i> , <i>kan</i> ^R	suicide vector	(5)
pSR47S- <i>P_{ahpc}</i> ::lux	R6Kori <i>sacB</i> , <i>kan</i> ^R <i>P_{ahpc}</i> ::lux	pSR47 containing <i>P. luminescens</i> lux operon	(6)
pJB3395	pBluescript:: <i>thyA</i> ⁺ <i>amp</i> ^R	<i>thyA</i> allelic exchange vector	J. Vogel
pTO243	pbluescript:: PolyHis- <i>attR1</i> -[<i>Kan</i> ^R - <i>Kan</i> ^R - <i>ccdB</i>]- <i>attR2</i>		O'Connor Tamara
pSK01	pSR47S:: $\Delta sdhA$	<i>sdhA</i> deletion plasmid	
pSK02	pSR47S:: $\Delta ridL$	<i>ridL</i> deletion plasmid	
pMMB207	<i>OriR</i> (RSF1010), Cm ^R		(7)
pMMB207 Δ 267	<i>OriR</i> (RSF1010), Cm ^R , Δ 267	pMMB207 lacking 267 bps of N-terminal <i>mobA</i>	Elizabeth Creasey
pSK03	pMMB207 Δ 267::PolyHis- <i>attR1</i> -[<i>Kan</i> ^R - <i>Kan</i> ^R - <i>ccdB</i>]- <i>attR2</i>	Gateway destination version of pMMB207 Δ 267	This work
pSK04	pMMB207 Δ 267::PolyHis- <i>attB1</i> - <i>sdeA</i> - <i>attB2</i>	<i>sdeA</i> complementation plasmid	This work
pSK05	pMMB207 Δ 267::PolyHis- <i>attB1</i> - <i>sdeB</i> - <i>attB2</i>	<i>sdeB</i> complementation plasmid	This work
pSK06	pMMB207 Δ 267::PolyHis- <i>attB1</i> - <i>sdeC</i> - <i>attB2</i>	<i>sdeC</i> complementation plasmid	This work
pSK07	pMMB207 Δ 267::PolyHis- <i>attB1</i> - <i>sdhA</i> - <i>attB2</i>	<i>sdhA</i> complementation plasmid	This work
pSdeB _{WT}	pJB908 PolyHis/c-myc- <i>attB1</i> - <i>sdeB</i> _{WT} - <i>attB2</i>	<i>sdeB</i> complementation plasmid	(8)
pSdeB _{C118S}	pJB908 PolyHis/c-myc- <i>attB1</i> - <i>sdeB</i> _{C118S} - <i>attB2</i>	pJB908 expressing inactive DUB domain of <i>sdeB</i>	Kristin Kotewicz
pSdeB _{H416A}	pJB908 PolyHis/c-myc- <i>attB1</i> - <i>sdeB</i> _{H416A} - <i>attB2</i>	pJB908 expressing inactive NP domain of <i>sdeB</i>	Kristin Kotewicz
pSdeB _{E859A}	pJB908 PolyHis/c-myc- <i>attB1</i> - <i>sdeB</i> _{E859A} - <i>attB2</i>	pJB908 expressing inactive ART domain of <i>sdeB</i>	(8)
pTO198	pSR47S:: $\Delta legA3$	<i>legA3</i> deletion plasmid	(9)

E. coli

DH5 α supE44 $\Delta lacU169(\Phi 80 lacZDM15)$

	<i>hsdR17 recA1 endA1 gyrA96</i> <i>thi-1 relA1</i>	
DH5α λpir	DH5α (<i>λpir</i>) <i>tet::Mu recA</i>	(10)
BL21 DE3	F ⁺ <i>ompT hsdSB dcm (DE3)</i>	

Oligonucleotides

Name	Sequences (5' to 3')
<i>Construction of sdhA mutant</i>	
SK1	GGCGCTAATTGCTGAAATCATTCAATATTAAAAAAATTAAC
SK2	CCGGGGGATGAACAATTACCCCTG
SK3	GATTCAGCAATTAGCGCCATCCGCATAAAATATTG
SK4	GAACTAGGGCGTAGGCGTTGACCATTAAAAG
pSR47s_sdhA_F	TTGTCATCCCCGGGCTGCAGGAAT
pSR47s_sdhA_R	CCTACGCCCTAGTTCTAGAGCGGCCGCC
<i>Construction of ridL mutant</i>	
SK5	TCATTATTATTATGTGTTCATTTAAGCCAAAAAAC
SK6	AGCCCAGGGGGTTATTACTGAAGTCGTGAC
SK7	CTAGAACTAGGATACTGGTGGATTGTCG
SK8	TGAACACATAATAATGACTTGGCTCTC
pSR47s_ridL_F	CAGTAATAACCCCCGGGCTGCAGGAAT
pSR47s_ridL_R	CACCAAGTATCCTAGTTCTAGAGCGGCCGCC
<i>Confirmation of recombinant plasmid</i>	
pSR47s_conF	GGGAACAAAAGCTGGAGC
pSR47s_conR	GTGAACGGCAGGTATATGT
<i>qRT-PCR</i>	
Name	Sequences (5' to 3')
rRNA_F	AGAGATGCATTAGTGCCTTCGGGA
rRNA_R	ACTAAGGATAAGGGTTGCGCTCGT
ridL_F	GTCCTCTGAAGGATAGCGAAAC
ridL_R	GTGTAAGTCCCGCAACAAATC
sidE_F	GCCTAAGTACGTTGAAGGGATAG
sidE_R	GCCTGTCAAGAGCACCTTA
sdeC_F	AAATCAGGAGAACCGGTTAGG
sdeC_R	CGTGAGAGCCGGATAATT
sdeB_F	CCAGGCTTCACTCACTTGATAA
sdeB_R	CCTCTCGATACCTACTGTGTCT
sdeA_F	CCCACTGCACCACAAGATAA
sdeA_R	GGTATACGGTTGCCAGATAG
sdhA_F	GGAAGGCAGGATTCTCCATTAA
sdhA_R	AGCTCTGAGTTCAGGAGGTAT
legA3_F	CTCCGCTCTTCAGATGAC
legA3_R	GAGTGGGTCGAGTGGGATAA

sidJ_F	GTTGTCCTACCCAACCTGG
sidJ_R	CAGAGAGGTGTCATGAGTGC

Mariner Tn-seq sequencing library construction

Name	Sequences (5' to 3')	Index
------	----------------------	-------

First PCR

Nextera 2A-R	GTCTCGTGGCTCGGAGATGTGTATAAGAGACAG
1st_TnR	GTAATACGACTCACTATAGGGCTAGAG

Second PCR- Leftward Mariner specific Nextera Indexed primers

mar147	AATGATACGGCGACCACCGAGATCTACACGCAGGCC GTTGACCGGGGACTTATCAGCCAACCTGTTA	GCAGGCCG
mar148	AATGATACGGCGACCACCGAGATCTACACAGGCAGAAC GTTGACCGGGGACTTATCAGCCAACCTGTTA	AGGCAGAA
mar149	AATGATACGGCGACCACCGAGATCTACACCCAGAGAGGC GTTGACCGGGGACTTATCAGCCAACCTGTTA	CAGAGAGG
mar150	AATGATACGGCGACCACCGAGATCTACACCGAGGCTGC GTTGACCGGGGACTTATCAGCCAACCTGTTA	CGAGGCTG
mar151	AATGATACGGCGACCACCGAGATCTACACAAGAGGCAC GTTGACCGGGGACTTATCAGCCAACCTGTTA	AAGAGGCA
mar152	AATGATACGGCGACCACCGAGATCTACACCGAGGAGCCC GTTGACCGGGGACTTATCAGCCAACCTGTTA	GAGGAGCC

Second PCR- Rightward Mariner specific Nextera Indexed primers

olk141	CAAGCAGAAGACGGCATACGAGATCCGCCTGCGTCTCGT GGGCTCGGAGATGTG	GCAGGCCG
N703 index	CAAGCAGAAGACGGCATACGAGATTCTGCCTGTCTCGT GGGCTCGGAGATGTG	AGGCAGAA

Reconditioning

P1	AATGATACGGCGACCACCGA
P2	CAAGCAGAAGACGGCATACGA

Sequencing

mar512	CGTTGACCGGGGACTTATCAGCCAACCTGTTA
--------	----------------------------------

SI References

1. K. H. Berger, R. R. Isberg, Two distinct defects in intracellular growth complemented by a single genetic locus in *Legionella pneumophila*. *Mol Microbiol* **7**, 7-19 (1993).
2. K. C. Jeong, J. A. Sexton, J. P. Vogel, Spatiotemporal regulation of a *Legionella pneumophila* T4SS substrate by the metaeffector SidJ. *PLoS Pathog* **11**, e1004695 (2015).
3. A. W. Ensminger, Y. Yassin, A. Miron, R. R. Isberg, Experimental evolution of *Legionella pneumophila* in mouse macrophages leads to strains with altered determinants of environmental survival. *PLoS Pathog* **8**, e1002731 (2012).
4. J. M. Park, S. Ghosh, T. J. O'Connor, Combinatorial selection in amoebal hosts drives the evolution of the human pathogen *Legionella pneumophila*. *Nat Microbiol* **5**, 599-609 (2020).
5. J. J. Merriam, R. Mathur, R. Maxfield-Boumil, R. R. Isberg, Analysis of the *Legionella pneumophila* fliI gene: intracellular growth of a defined mutant defective for flagellum biosynthesis. *Infect Immun* **65**, 2497-2501 (1997).
6. J. Coers, R. E. Vance, M. F. Fontana, W. F. Dietrich, Restriction of *Legionella pneumophila* growth in macrophages requires the concerted action of cytokine and Naip5/Ipfaf signalling pathways. *Cell Microbiol* **9**, 2344-2357 (2007).
7. V. M. Morales, A. Backman, M. Bagdasarian, A series of wide-host-range low-copy-number vectors that allow direct screening for recombinants. *Gene* **97**, 39-47 (1991).
8. K. M. Kotewicz *et al.*, A Single *Legionella* Effector Catalyzes a Multistep Ubiquitination Pathway to Rearrange Tubular Endoplasmic Reticulum for Replication. *Cell Host Microbe* **21**, 169-181 (2017).
9. T. J. O'Connor, D. Boyd, M. S. Dorer, R. R. Isberg, Aggravating genetic interactions allow a solution to redundancy in a bacterial pathogen. *Science* **338**, 1440-1444 (2012).
10. R. Kolter, M. Inuzuka, D. R. Helinski, Trans-complementation-dependent replication of a low molecular weight origin fragment from plasmid R6K. *Cell* **15**, 1199-1208 (1978).