
## **Supplemental Online Content**



eFigure. Flow diagram of the study cohort

Patients with medically managed missed abortion between 1/1/2016 and 12/31/2020 (ICD-10 codes: O02.0 and O02.1) Patients excluded: - individuals outside of age requirement of 15 to 49 - individuals had both a missed abortion and at least one exclusion diagnosis occurring within 60 days prior to their missed abortion (exclusion dx: Incomplete abortion: O03.4, O03.3x; complete or unspecified abortion: O03.9; stillbirth: P95; elective abortion: Z33.2) - individuals had an exclusion diagnosis (O03.4, O03.3x, O03.9, P95, Z33.2) on the same day of filling a prescription for either mifepristone or misoprostol, or had an exclusion diagnosis following a missed abortion diagnosis but before the medication was dispensed - individuals with procedural codes indicative of surgical completion of missed abortion (59820 or 59821) on the same day or within 60 days prior to an individual's outpatient services claims - individuals received mifepristone but never received misoprostol Final cohort, established using both prescription drug claims and outpatient services claims (S0191, S0190) **Combination Therapy:** Patients Monotherapy: Patients who who received combination therapy received monotherapy (misoprostol (misoprostol & mifepristone) within only) within 28 days of a missed 28 days of a missed abortion dx abortion dx