

**S2 Table: Cell population distribution of isolated PMN from HITTIN and HIT, as determined by flow cytometry**

Characteristic	HITTIN <sup>a</sup> , N = 14 (3 missing values) <sup>*</sup>	HIT <sup>b</sup> , N = 10 (1 missing value) <sup>*</sup>	p-value <sup>**</sup>
<b>CD15+ CD66b+</b>	92.70% (87.05, 93.53)	94.35% (93.65, 97.08)	0.02**
<b>CD16+ (Neutrophils)</b>	86.90 (80.28, 92.90)	90.45 (86.10, 93.25)	0.28
<b>CD16- CD14<sub>low</sub> (Eosinophils)</b>	4.33 (1.51, 7.23)	3.63 (2.11, 6.60)	0.93
<b>CD15- CD66b-</b>	7.31% (6.44, 12.95)	5.67% (2.92, 6.32)	0.03**
<b>CD3- CD14- (Other)</b>	0.30 (0.17, 0.74)	0.39 (0.18, 0.58)	0.66
<b>CD3- CD14+ (Monocytes)</b>	0.07 (0.04, 0.12)	0.11 (0.07, 0.19)	0.22
<b>CD3+ (T-cells)</b>	6.70 (6.10, 12.43)	4.55 (2.27, 6.00)	0.02**

\*Median (IQR); \*\*Wilcoxon rank sum test

<sup>a</sup>HITTIN (HIV-1-infected persistently TB, tuberculin and IGRA negative), <sup>b</sup>HIT (HIV-1-infected IGRA positive tuberculin positive)