

Supplement Table S1. Participant Characteristics.

Characteristic	N = 32
Age, mean (SD); range	56 (12); 21-74
Sex, n (%)	
Male	14 (44)
Female	18 (56)
Identified as Hispanic	15 (53)
Dialysis modality in-center hemodialysis, n (%)	32 (100)
Vascular access	
Arteriovenous fistula	18 (56)
Arteriovenous graft	7 (22)
Central venous catheter	7 (22)
Prior kidney transplant	4 (13)
Unemployed, n (%)	30 (94)
Education level, n (%)	
No School	0 (0)
Less than high-school	25 (78)
High-school	2 (6)
More than high-school	5 (16)
Health insurance, n (%)	
Medicare	1 (3)
Medicaid	9 (28)
Medicare and Medicaid	21 (66)
Other	1 (3)
Annual income	
<\$25,000	27 (84)
\$25,000-\$50,000	5 (16)
Difficulty affording medications, n (%)	8 (25)
Housing, n (%)	14 (44)
Food insecurity, n (%)	20 (63)
Utility issues, n (%)	8 (25)
Transportation issues, n (%)	17 (53)
Comorbidities, n (%)	
Myocardial infarction	5 (16%)
Heart failure	11 (34%)
Stroke	8 (25%)
Heart valve problem	5 (16%)
Coronary bypass	1 (3%)
Hypertension	31 (97%)
Lung disease	7 (22%)
Diabetes	19 (59%)
Amputation	4 (13%)
Peptic ulcer disease	9 (28%)
Liver disease	1 (3%)
Anemia	15 (47%)
Cancer	5 (16%)