

Supplementary Materials

Phase 3 Clinical Trial Comparing the Safety and Efficacy of Netarsudil to Ripasudil in Patients with Primary Open-Angle Glaucoma or Ocular Hypertension: Japan Rho Kinase Elevated Intraocular Pressure Treatment Trial (J-ROCKET)

Makoto Araie¹; Kazuhisa Sugiyama²; Kenji Aso³; Koji Kanemoto³; Ryo Iwata³; David A. Hollander⁴; Michelle Senchyna⁴; Casey C. Kocczynski⁴

¹Sekikawa Hospital, Tokyo, Japan

²Department of Ophthalmology, Kanazawa University Graduate School of Medical Science, Ishikawa, Japan

³Aerie Pharmaceuticals, Ireland Ltd. (Japan Branch), Tokyo, Japan

⁴Aerie Pharmaceuticals, Inc., Durham, NC., USA

Corresponding Author:

Makoto Araie

Sekikawa Hospital, 1-4-1, Nishinippori, Arakawa-ku, Tokyo

e-mail: m.araie-tyk@nifty.com

Institutional Review Board (IRB) Information

Shin-Akasaka Clinic IRB

5-5-1, Roppongi, Minato-ku, Tokyo, 106-0032, Japan

Telephone: +81-3-5770-1250

Yokohama Minoru Clinic IRB

1-13-8, Bessho, Minami-ku, Yokohama-shi, Kanagawa, 232-0064, Japan

Telephone: +81-45-722-8655

Japan Conference of Clinical Research IRB

1-13-23, Minamiikebukuro, Toshima-ku, Tokyo, 171-0022, Japan

Telephone: +81-3-6868-7022

Adachi Kyosai Hospital IRB

1-36-8, Yanagihara, Adachi-ku, Tokyo, 120-0022, Japan

Telephone: +81-3-3881-6116

Kobori Central IRB

5-11-8, Nishinippori, Arakawa-ku, Tokyo, 116-0013, Japan

Telephone: +81-3-3805-8181

Osaka Pharmacology Clinical Research Hospital IRB

4-1-29, Miyahara, Yodogawa-ku, Osaka-shi, Osaka, 532-0003, Japan

Telephone: +81-6-6395-9000