Supplementary Table S11. Renal cancer patient characteristics, median survival, and Multivariate Cox Regression Modeling.

| | x Regression i | | | | | | |
|---|---|-------------------|----------------------------|---------------------|-------------|-------------|---------|
| Patient Characte | | LOR | ALP | | No Benzo | | P-value |
| _ | N | 73 (12.4) | | 38 (6.5) | 477 (81.1) | 588 (100%) | |
| Sex | Female | 25 (34.2%) | 21 (55.3%) | | 185 (38.8%) | 231 (39.3%) | 0.087 |
| | Male | 48 (65.8%) | · · | 44.7%) | 292 (61.2%) | 357 (60.7%) | |
| Grade (Clinical) | <u> </u> | | | (7.9%) | 23 (4.9%) | 26 (4.5%) | 0.288 |
| | II | 18 (24.7%) | | | 131 (28.0%) | 160 (27.6%) | |
| | III | 24 (32.9%) | | 34.2%) | 135 (28.8%) | 172 (29.7%) | |
| | IV | 6 (8.2%) | | (5.3%) | 59 (12.6%) | 67 (11.6%) | |
| | Not Reported | 25 (34.2%) | | 23.7%) | 120 (25.6%) | 154 (26.6%) | |
| Grade (Pathological) | I | | | (7.9%) | 23 (4.9%) | 26 (4.5%) | 0.435 |
| | | 18 (24.7%) | | 26.3%) | 129 (27.5%) | 157 (27.1%) | |
| | III | 23 (31.5%) | | 34.2%) | 136 (29.0%) | 172 (29.7%) | |
| | IV | 8 (11.0%) | | (5.3%) | 60 (12.8%) | 70 (12.1%) | |
| | Not Reported | 24 (32.9%) | 10 (| 26.3%) | 121 (25.8%) | 155 (26.7%) | |
| Stage (Clinical) | I | 13 (17.8%) | 9 (| 24.3%) | 133 (27.9%) | 155 (26.4%) | 0.412 |
| | | 4 (5.5%) | 1 (2.7%) | | 44 (9.2%) | 49 (8.3%) | |
| | III | 4 (5.5%) | 1 (2.7%) | | 24 (5.0%) | 29 (4.9%) | |
| | IV | 16 (21.9%) | 7 (| 18.9%) | 78 (16.4%) | 101 (17.2%) | |
| | Not Reported | 36 (49.3%) | 19 (| 51.4%) | 198 (41.5%) | 253 (43.1%) | |
| Stage (Pathological) | I | 10 (13.7%) | 7 (| 18.9%) | 106 (22.2%) | 123 (21.0%) | 0.482 |
| | Ш | 3 (4.1%) | 2 | (5.4%) | 32 (6.7%) | 37 (6.3%) | |
| | III | 5 (6.8%) | 2 | (5.4%) | 47 (9.9%) | 54 (9.2%) | |
| | IV | 9 (12.3%) | 5 (| 13.5%) | 39 (8.2%) | 53 (9.0%) | |
| | Not Reported | 46 (63.0%) | 21 (| 56.8%) | 253 (53.0%) | 320 (54.5%) | |
| Overall Survival Indicator | Alive | 26 (35.6%) | 26 (35.6%) 16 (| | 229 (48.0%) | 271 (46.1%) | 0.124 |
| | Dead | 47 (64.4%) 22 (| | 57.9%) | 248 (52.0%) | 317 (53.9%) | |
| Progression-Free Survival Indicator | No Progression | 20 (27.4%) | 20 (27.4%) 13 (| | 205 (43.0%) | 238 (40.5%) | 0.030 |
| | Progression | 53 (72.6%) | 25 (| 65.8%) | 272 (57.0%) | 350 (59.5%) | |
| Overall Survival (OS) | Median Surv. (95% Cl) | Sample: K | Log Rank Kidney P-value | | | | |
| Total | 82.4 (65.6, 96.0) | E=317 C=271 T=588 | | p= 0.272 | | | |
| ALP | 66.9 (24.0, 120.6) | E=22 C=16 T=38 | | | | | |
| LOR | 66.9 (48.3, 96.0) | E=47 C=26 T=73 | | | - | | |
| No Benzo | 86.8 (66.2, 105.9) | E=248 C=229 T=477 | | | _ | | |
| Progression-Free Survival (PFS) | Median Surv. (95% CI) | Sample: Kidney | | Log Rank P-value | | | |
| Total | 49.1 (38.3, 66.8) | E=350 C=23 | - | p= 0.051 | | | |
| ALP | 35.6 (13.9, 95.7) | E=25 C=13 T=38 | | | - | | |
| LOR | 36.8 (26.1, 53.8) | E=53 C=20 T=73 | | | - | | |
| No Benzo | 59.8 (39.1, 79.2) | E=272 C=205 T=477 | | | - | | |
| | | 2 212 0 20 | •••• | | | | |
| Overall Survival (C Predictors: Cohort |)S) t, Sex, Clinical Grad | e, and Clinical | Stage | | | | |
| Cohort | HR (95% CI) | | P-value | | | | |
| No Benzo | Ref. | | 0.6198 | | | | |
| ALP | 1.153 (0.741 – | 1.795) | 0.5268 | | | | |
| LOR | 0.898 (0.651 – | | 0.5132 | | | | |
| Progression-Free | | , | | | | | |
| No Benzo | Ref. | | 0.7907 | | | | |
| ALP | 1.152 (0.760 – 1.747) | | 0.5058 | | | | |
| LOR | | | 0.8034 | | | | |
| LOIN | 1.039 (0.768 – 1.407) ensored T= Total | | 0.0004 | | | | |

E= event, C= censored, T= Total