

**Supplemental Appendix 4. Summary of AMR deaths and DALYs by infectious syndrome for the WHO Region of the Americas countries, 2019**

Country	Infectious Syndrome	Attributable DALYs Counts	Age standardised rate per 100k	Associated DALYs Counts	Age standardised rate per 100k	Attributable Death Counts	Age standardised rate per 100k	Associated Death Counts	Age standardised rate per 100k
Antigua and Barbuda	All	49 (33-68)	55.6 (37.9-77.1)	1,150 (755-1,640)	1303.6 (862.3-1859.9)	12 (8-18)	14.2 (9.5-20.2)	291 (187-432)	329.2 (212-482.1)
Antigua and Barbuda	Bacterial skin infections	4 (2-6)	4.2 (2.4-7.1)	73 (42-126)	78 (45.6-133.1)	1 (0-1)	0.9 (0.5-1.6)	17 (9-29)	17.7 (9.7-30.2)
Antigua and Barbuda	Bone and joint infections	0 (0-0)	0.2 (0.1-0.6)	4 (1-10)	4.5 (1.3-10.6)	0 (0-0)	0.1 (0-0.1)	1 (0-3)	1.2 (0.3-2.8)
Antigua and Barbuda	BSI	17 (9-28)	18.6 (9.8-31.6)	407 (216-656)	465.5 (251.7-744)	4 (2-7)	5 (2.6-8.4)	108 (57-176)	122.9 (66.6-199.9)
Antigua and Barbuda	Cardiac infections	1 (1-1)	1 (0.7-1.3)	26 (19-35)	26.2 (19.4-35.2)	0 (0-0)	0.3 (0.2-0.4)	7 (5-9)	7 (5-9.5)
Antigua and Barbuda	CNS infections	0 (0-1)	0.5 (0.3-0.8)	18 (12-28)	23.9 (15.4-36.8)	0 (0-0)	0.1 (0.1-0.2)	4 (3-7)	5.5 (3.4-8.9)
Antigua and Barbuda	Diarrhoea	0 (0-0)	0 (0-0)	1 (1-2)	1.6 (0.8-2.8)	0 (0-0)	0 (0-0)	0 (0-0)	0.3 (0.2-0.5)
Antigua and Barbuda	Gonorrhoea and chlamydia			1 (0-1)	0.8 (0.4-1.6)			0 (0-0)	0.1 (0-0.2)
Antigua and Barbuda	Intra-abdominal infections	8 (5-13)	8.8 (5.6-13.3)	198 (122-305)	199.3 (122.6-306.9)	2 (1-3)	2.2 (1.4-3.4)	50 (30-81)	50.8 (29.9-81.1)
Antigua and Barbuda	LRI and thorax infections	14 (10-20)	17.3 (12.7-23.4)	335 (233-465)	414.2 (288.7-574.8)	4 (3-5)	4.4 (3.1-6)	83 (56-120)	102 (69.1-145.4)
Antigua and Barbuda	TB	0 (0-0)	0 (0-0)	0 (0-1)	0.4 (0.1-1.3)	0 (0-0)	0 (0-0)	0 (0-1)	0.1 (0-0.5)
Antigua and Barbuda	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0.1 (0-0.1)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Antigua and Barbuda	UTI	4 (3-6)	4.9 (3.5-6.9)	85 (59-120)	89.1 (62.7-125.3)	1 (1-2)	1.2 (0.8-1.7)	21 (14-30)	21.7 (14.9-31.7)

Argentina	All	35,300 (27,400-44,700)	65.1 (50.4-82.6)	701,000 (525,000-917,000)	1413.3 (1049.9-1851.1)	8,980 (6,780-11,800)	16.6 (12.5-21.8)	178,000 (131,000-240,000)	358.7 (262.3-486.5)
Argentina	Bacterial skin infections	1,080 (618-1,830)	2 (1.1-3.3)	18,500 (10,200-32,400)	34.9 (19.3-60.4)	259 (139-447)	0.5 (0.3-0.8)	4,420 (2,350-7,800)	8.4 (4.4-14.7)
Argentina	Bone and joint infections	68 (18-160)	0.1 (0-0.3)	1,370 (371-3,290)	2.6 (0.7-6.3)	18 (5-43)	0 (0-0.1)	356 (98-857)	0.7 (0.2-1.7)
Argentina	BSI	7,820 (4,260-13,100)	14.8 (8.2-24.7)	188,000 (107,000-298,000)	400.7 (236.9-621.6)	2,050 (1,090-3,430)	3.9 (2.1-6.5)	49,100 (27,400-78,700)	104.9 (60.5-165.3)
Argentina	Cardiac infections	849 (686-1,040)	1.5 (1.3-1.9)	16,400 (13,600-20,700)	31.4 (26-39.8)	223 (172-278)	0.4 (0.3-0.5)	4,280 (3,400-5,450)	8.2 (6.5-10.5)
Argentina	CNS infections	168 (126-238)	0.3 (0.3-0.5)	6,860 (5,060-9,560)	15.9 (11.6-22.2)	39 (29-57)	0.1 (0.1-0.1)	1,590 (1,140-2,280)	3.7 (2.6-5.3)
Argentina	Diarrhoea	22 (11-39)	0 (0-0.1)	1,440 (757-2,430)	3.5 (1.9-6.2)	5 (2-9)	0 (0-0)	278 (142-484)	0.7 (0.3-1.2)
Argentina	Gonorrhoea and chlamydia			262 (147-432)	0.6 (0.3-0.9)			26 (8-51)	0.1 (0-0.1)
Argentina	Intra-abdominal infections	5,600 (3,750-7,930)	10.3 (6.9-14.7)	118,000 (76,100-170,000)	226.6 (145.6-329.4)	1,440 (960-2,030)	2.7 (1.8-3.8)	30,200 (19,500-43,800)	58.3 (37.4-84.9)
Argentina	LRI and thorax infections	16,100 (13,700-18,900)	29.4 (25.1-34.9)	294,000 (247,000-358,000)	590.7 (491.8-725)	4,080 (3,170-5,170)	7.5 (5.8-9.5)	73,800 (57,400-94,300)	147.5 (114.6-189.8)
Argentina	TB	48 (21-118)	0.1 (0-0.2)	1,650 (734-4,020)	3.4 (1.5-8.3)	20 (3-66)	0 (0-0.1)	631 (103-2,140)	1.3 (0.2-4.4)
Argentina	Typhoid, paratyphoid, and iNTS	0 (0-1)	0 (0-0)	13 (2-32)	0 (0-0.1)	0 (0-0)	0 (0-0)	3 (0-8)	0 (0-0)
Argentina	UTI	3,570 (2,500-4,580)	6.4 (4.5-8.2)	55,200 (38,800-73,800)	102.9 (72.5-138)	859 (599-1,140)	1.5 (1.1-2.1)	13,400 (9,150-18,500)	24.9 (17.2-34.7)
Barbados	All	312 (226-419)	67.6 (48.6-91.2)	6,040 (4,160-8,420)	1563.3 (1067.4-2210.7)	79 (55-109)	17 (11.9-23.7)	1,520 (1,020-2,160)	392.1 (259.7-562.7)
Barbados	Bacterial skin infections	35 (22-58)	7.4 (4.7-12.1)	600 (372-982)	136.8 (85.3-221)	8 (4-14)	1.7 (0.9-2.9)	135 (78-229)	31.1 (18.2-51.6)
Barbados	Bone and joint infections	1 (0-2)	0.2 (0.1-0.5)	18 (5-44)	4.2 (1.1-10)	0 (0-1)	0.1 (0-0.1)	5 (1-11)	1.1 (0.3-2.6)
Barbados	BSI	86 (44-144)	19.3 (10.1-31.9)	1,910 (1,000-3,080)	551 (312.4-860.9)	22 (11-38)	5 (2.6-8.3)	488 (257-807)	140.5 (78-224.4)
Barbados	Cardiac infections	6 (4-7)	1.2 (0.9-1.6)	135 (100-179)	33.1 (24.7-44)	1 (1-2)	0.3 (0.2-0.4)	35 (25-47)	8.5 (6.2-11.5)

Barbados	CNS infections	2 (1-3)	0.4 (0.3-0.7)	56 (36-93)	20.6 (13.1-33.5)	0 (0-1)	0.1 (0.1-0.2)	13 (8-22)	4.8 (3-8)
Barbados	Diarrhoea	0 (0-0)	0 (0-0)	5 (2-10)	1.6 (0.8-3.2)	0 (0-0)	0 (0-0)	1 (0-1)	0.3 (0.1-0.5)
Barbados	Gonorrhoea and chlamydia			2 (1-5)	0.9 (0.4-1.8)			0 (0-1)	0.1 (0-0.2)
Barbados	Intra-abdominal infections	40 (25-62)	8.4 (5.2-13.1)	854 (514-1,350)	193.3 (113.9-309.4)	10 (6-16)	2.1 (1.3-3.4)	219 (129-349)	49.6 (28.8-80.9)
Barbados	LRI and thorax infections	97 (74-128)	21.2 (16.1-28)	1,700 (1,220-2,350)	455 (319-635.2)	25 (18-34)	5.5 (3.9-7.5)	438 (303-618)	115.6 (77-165.5)
Barbados	TB	0 (0-0)	0 (0-0)	1 (0-3)	0.3 (0.1-0.9)	0 (0-0)	0 (0-0)	0 (0-1)	0.1 (0-0.4)
Barbados	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	1 (0-1)	0.2 (0.1-0.4)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0.1)
Barbados	UTI	45 (32-57)	9.3 (6.6-11.8)	758 (541-973)	166.3 (119.7-214.2)	11 (8-14)	2.3 (1.6-3)	185 (130-245)	40.6 (28.5-53.4)
Belize	All	158 (108-220)	57.8 (39.8-79.9)	5,310 (3,630-7,450)	1616.2 (1108.9-2259.8)	39 (27-56)	14.5 (9.8-20.4)	1,310 (873-1,870)	399.8 (268.7-567.2)
Belize	Bacterial skin infections	6 (3-12)	2.4 (1-4.7)	152 (70-295)	50.3 (22.1-97.3)	1 (1-3)	0.5 (0.2-1.1)	34 (15-67)	11.1 (4.6-21.9)
Belize	Bone and joint infections	1 (0-2)	0.3 (0.1-0.6)	17 (5-42)	5.5 (1.5-13.1)	0 (0-0)	0.1 (0-0.1)	4 (1-11)	1.4 (0.4-3.4)
Belize	BSI	49 (28-79)	17.8 (9.7-28.9)	1,800 (1,100-2,730)	542.4 (324.3-827.6)	13 (7-21)	4.6 (2.5-7.6)	463 (277-713)	139.5 (82.3-216.2)
Belize	Cardiac infections	1 (1-2)	0.4 (0.3-0.6)	46 (32-64)	13 (9.3-18.1)	0 (0-0)	0.1 (0.1-0.2)	12 (8-17)	3.4 (2.4-4.8)
Belize	CNS infections	2 (1-3)	0.4 (0.2-0.8)	89 (52-159)	23 (13.6-41)	0 (0-1)	0.1 (0.1-0.2)	20 (12-37)	5.2 (3-9.6)
Belize	Diarrhoea	0 (0-0)	0 (0-0.1)	9 (5-15)	2.5 (1.4-4.3)	0 (0-0)	0 (0-0)	2 (1-3)	0.5 (0.3-0.9)
Belize	Gonorrhoea and chlamydia			3 (1-6)	0.7 (0.3-1.3)			0 (0-1)	0.1 (0-0.2)
Belize	Intra-abdominal infections	31 (20-46)	11.3 (7.2-16.6)	924 (554-1,400)	288.5 (177.9-429.3)	8 (5-12)	2.8 (1.8-4.3)	232 (136-364)	72.4 (43.5-112.7)
Belize	LRI and thorax infections	54 (40-71)	19.8 (14.8-26.3)	1,930 (1,440-2,570)	578 (431.8-772.6)	13 (10-18)	4.9 (3.5-6.7)	459 (329-623)	138.9 (99.9-188.1)
Belize	TB	0 (0-1)	0.1 (0-0.3)	11 (4-34)	3 (1.1-9)	0 (0-0)	0 (0-0.1)	5 (1-19)	1.3 (0.2-5.1)

Belize	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-1)	0.1 (0-0.1)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Belize	UTI	13 (10-19)	5.3 (3.9-7.3)	327 (241-452)	109.3 (80-151.8)	3 (2-5)	1.3 (0.9-1.8)	78 (55-111)	26 (18.3-37.3)
Bolivia	All	10,100 (7,220-13,600)	116.9 (83.7-158)	373,000 (272,000-508,000)	3232.9 (2339.8-4406.2)	2,490 (1,770-3,430)	29 (20.4-40.2)	90,500 (64,300-123,000)	791.2 (562.1-1088.2)
Bolivia	Bacterial skin infections	141 (44-324)	1.9 (0.6-4.2)	3,300 (1,250-7,060)	36.1 (12.8-79.7)	31 (9-72)	0.4 (0.1-0.9)	709 (255-1,560)	7.8 (2.6-17.6)
Bolivia	Bone and joint infections	22 (6-51)	0.3 (0.1-0.6)	514 (132-1,240)	5.5 (1.4-13.2)	5 (1-12)	0.1 (0-0.2)	120 (30-303)	1.3 (0.3-3.2)
Bolivia	BSI	2,430 (1,440-3,770)	26 (14.8-41.9)	108,000 (69,500-155,000)	865.9 (549.8-1281.9)	594 (352-941)	6.4 (3.6-10.4)	26,300 (16,900-38,300)	211.3 (131.8-316.3)
Bolivia	Cardiac infections	55 (39-75)	0.6 (0.4-0.9)	1,810 (1,250-2,520)	17.1 (12-23.5)	13 (9-18)	0.2 (0.1-0.2)	438 (300-611)	4.1 (2.8-5.7)
Bolivia	CNS infections	70 (44-113)	0.6 (0.4-1)	4,070 (2,560-6,400)	31.1 (19.8-49.4)	16 (10-26)	0.1 (0.1-0.2)	906 (562-1,470)	6.9 (4.3-11.2)
Bolivia	Diarrhoea	7 (3-14)	0.1 (0-0.1)	656 (271-1,350)	4.8 (2-9.6)	2 (1-4)	0 (0-0)	129 (46-295)	0.9 (0.3-2.1)
Bolivia	Gonorrhoea and chlamydia			69 (29-149)	0.6 (0.2-1.2)			7 (1-18)	0.1 (0-0.2)
Bolivia	Intra-abdominal infections	1,850 (1,160-2,680)	22.3 (14.2-32.1)	48,000 (29,600-71,900)	499.5 (312.2-735.3)	456 (286-687)	5.5 (3.5-8.3)	11,800 (7,030-18,200)	123.1 (74.6-187.4)
Bolivia	LRI and thorax infections	4,780 (3,570-6,210)	56.2 (41.8-72.7)	187,000 (140,000-240,000)	1571.4 (1182.1-2033.6)	1,150 (814-1,530)	13.6 (9.6-18.3)	43,600 (31,200-58,000)	370.6 (267.2-492.7)
Bolivia	TB	244 (89-615)	2.6 (0.9-6.7)	9,240 (3,380-23,700)	85.5 (31.2-216.1)	114 (15-388)	1.2 (0.2-4.2)	4,050 (533-13,800)	37.6 (5-127.9)
Bolivia	Typhoid, paratyphoid, and iNTS	1 (0-2)	0 (0-0)	51 (16-118)	0.4 (0.1-0.9)	0 (0-1)	0 (0-0)	10 (2-28)	0.1 (0-0.2)
Bolivia	UTI	465 (297-712)	6.3 (4.1-9.5)	10,300 (6,420-16,000)	115.1 (72.7-176.7)	111 (69-170)	1.5 (0.9-2.3)	2,460 (1,520-3,870)	27.4 (16.9-42.4)
Brazil	All	138,000 (103,000-182,000)	63.2 (47.4-82.9)	3,570,000 (2,630,000-4,810,000)	1724.4 (1267.8-2306)	33,200 (24,200-45,000)	15.2 (11.2-20.4)	859,000 (621,000-1,180,000)	413.5 (299.7-569.1)
Brazil	Bacterial skin infections	4,910 (2,490-8,660)	2.2 (1.1-3.9)	107,000 (54,800-183,000)	47.1 (24.5-80.7)	995 (480-1,820)	0.4 (0.2-0.8)	21,400 (10,300-38,400)	9.5 (4.6-16.7)

Brazil	Bone and joint infections	393 (109-927)	0.2 (0-0.4)	9,320 (2,610-22,300)	4.1 (1.2-9.7)	87 (24-206)	0 (0-0.1)	2,060 (564-4,980)	0.9 (0.2-2.2)
Brazil	BSI	34,000 (19,000-55,700)	16 (9.2-25.8)	1,080,000 (665,000-1,640,000)	558.6 (354-830)	8,520 (4,830-13,900)	4 (2.3-6.5)	269,000 (166,000-411,000)	139.3 (87.7-210.9)
Brazil	Cardiac infections	1,990 (1,490-2,500)	0.9 (0.7-1.1)	55,700 (43,700-69,500)	24.4 (19.2-30.5)	498 (373-632)	0.2 (0.2-0.3)	13,900 (10,800-17,700)	6.1 (4.7-7.8)
Brazil	CNS infections	785 (565-1,170)	0.4 (0.3-0.6)	43,200 (31,000-61,000)	22.9 (16.2-32.2)	176 (125-266)	0.1 (0.1-0.1)	9,580 (6,800-13,800)	5.1 (3.5-7.2)
Brazil	Diarrhoea	87 (54-131)	0 (0-0.1)	8,030 (4,590-13,700)	4.7 (2.7-7.9)	18 (11-30)	0 (0-0)	1,410 (787-2,330)	0.8 (0.5-1.4)
Brazil	Gonorrhoea and chlamydia			1,230 (699-1,910)	0.5 (0.3-0.8)			120 (31-242)	0.1 (0-0.1)
Brazil	Intra-abdominal infections	25,200 (17,100-35,600)	10.9 (7.5-15.4)	628,000 (413,000-909,000)	268.4 (176.8-390.5)	6,390 (4,310-8,950)	2.8 (1.9-3.9)	159,000 (104,000-231,000)	68 (44.3-98.3)
Brazil	LRI and thorax infections	53,500 (43,500-67,100)	24.8 (20.2-31.1)	1,320,000 (1,050,000-1,680,000)	649.4 (509-826.6)	12,500 (9,480-16,100)	5.8 (4.4-7.4)	303,000 (228,000-394,000)	148.7 (111.3-194.3)
Brazil	TB	596 (264-1,270)	0.3 (0.1-0.5)	22,200 (10,100-46,600)	9.4 (4.3-19.8)	260 (41-799)	0.1 (0-0.3)	8,820 (1,420-26,900)	3.7 (0.6-11.5)
Brazil	Typhoid, paratyphoid, and iNTS	7 (3-13)	0 (0-0)	343 (132-717)	0.2 (0.1-0.4)	1 (0-3)	0 (0-0)	70 (17-170)	0 (0-0.1)
Brazil	UTI	16,500 (12,000-20,200)	7.5 (5.5-9.2)	303,000 (230,000-382,000)	134.7 (102.7-169.8)	3,820 (2,790-4,880)	1.7 (1.3-2.2)	70,400 (52,300-92,000)	31.3 (23.3-40.6)
Canada	All	12,900 (8,830-18,100)	18.3 (12.3-25.9)	229,000 (151,000-328,000)	395.6 (256.2-576.5)	2,890 (1,960-4,180)	4.1 (2.7-6)	51,000 (33,100-75,800)	88 (55.4-132.9)
Canada	Bacterial skin infections	357 (158-721)	0.5 (0.2-1)	6,140 (2,730-12,700)	10.2 (4.6-21)	69 (28-149)	0.1 (0-0.2)	1,170 (489-2,520)	1.9 (0.8-4.2)
Canada	Bone and joint infections	52 (17-118)	0.1 (0-0.2)	961 (308-2,150)	1.6 (0.5-3.6)	11 (3-26)	0 (0-0)	195 (61-466)	0.3 (0.1-0.8)
Canada	BSI	4,660 (2,480-7,840)	6.9 (3.7-11.5)	93,300 (50,300-152,000)	171.6 (92.2-279.1)	1,060 (550-1,810)	1.6 (0.8-2.7)	21,200 (11,300-35,600)	38.8 (20.9-64.4)
Canada	Cardiac infections	456 (268-646)	0.6 (0.4-0.9)	8,170 (4,980-11,700)	13.7 (8.4-19.7)	104 (60-151)	0.1 (0.1-0.2)	1,830 (1,100-2,690)	3.1 (1.8-4.5)
Canada	CNS infections	35 (21-67)	0.1 (0-0.1)	1,250 (764-2,200)	3.6 (2.3-5.9)	7 (4-13)	0 (0-0)	229 (134-412)	0.7 (0.4-1.1)
Canada	Diarrhoea	19 (10-32)	0 (0-0)	765 (384-1,350)	1.9 (1-3.4)	4 (2-7)	0 (0-0)	120 (63-200)	0.3 (0.1-0.5)

Canada	Gonorrhoea and chlamydia			58 (32-100)	0.2 (0.1-0.3)			6 (2-12)	0 (0-0)
Canada	Intra-abdominal infections	3,020 (1,930-4,480)	4.3 (2.7-6.4)	57,100 (35,900-84,400)	95.3 (58.9-144.5)	701 (446-1,040)	1 (0.6-1.5)	13,300 (8,340-19,800)	22.2 (13.7-33.4)
Canada	LRI and thorax infections	2,990 (2,320-3,830)	4 (3.1-5.2)	42,600 (31,600-58,000)	70 (51.4-96.5)	640 (460-880)	0.8 (0.6-1.2)	9,110 (6,320-13,000)	14.9 (10.3-21.3)
Canada	TB	15 (10-24)	0 (0-0)	312 (205-468)	0.5 (0.4-0.8)	5 (0-11)	0 (0-0)	79 (9-193)	0.1 (0-0.3)
Canada	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	4 (1-8)	0 (0-0)	0 (0-0)	0 (0-0)	1 (0-2)	0 (0-0)
Canada	UTI	1,330 (1,040-1,720)	1.7 (1.3-2.2)	18,000 (13,500-24,500)	27 (20-37.3)	284 (216-378)	0.4 (0.3-0.5)	3,840 (2,790-5,410)	5.8 (4.1-8.2)
Chile	All	10,200 (7,360-13,700)	43.7 (31.6-59.1)	201,000 (138,000-282,000)	930.9 (630.6-1312)	2,720 (1,930-3,770)	11.7 (8.3-16.2)	53,800 (36,700-77,100)	247.6 (167.4-357.6)
Chile	Bacterial skin infections	273 (140-500)	1.2 (0.6-2.1)	4,700 (2,390-9,090)	20.5 (10.5-39.4)	66 (32-125)	0.3 (0.1-0.5)	1,130 (540-2,220)	4.9 (2.4-9.7)
Chile	Bone and joint infections	26 (8-62)	0.1 (0-0.3)	568 (167-1,320)	2.5 (0.7-5.8)	7 (2-17)	0 (0-0.1)	151 (45-363)	0.7 (0.2-1.6)
Chile	BSI	2,810 (1,510-4,700)	12.2 (6.6-20.4)	62,800 (34,500-101,000)	305.9 (169.8-485.6)	766 (398-1,300)	3.3 (1.7-5.6)	17,000 (9,330-27,900)	82.8 (45.5-135.6)
Chile	Cardiac infections	218 (154-311)	0.9 (0.7-1.3)	4,730 (3,460-6,620)	20.8 (15.2-29.3)	60 (41-87)	0.3 (0.2-0.4)	1,290 (908-1,850)	5.6 (4-8.1)
Chile	CNS infections	52 (37-79)	0.3 (0.2-0.4)	2,100 (1,530-3,070)	11.9 (8.5-17.4)	12 (9-19)	0.1 (0-0.1)	488 (344-734)	2.8 (1.9-4.1)
Chile	Diarrhoea	8 (4-13)	0 (0-0.1)	381 (218-616)	2.2 (1.2-3.6)	2 (1-3)	0 (0-0)	72 (37-123)	0.4 (0.2-0.7)
Chile	Gonorrhoea and chlamydia			82 (43-168)	0.4 (0.2-0.9)			8 (2-19)	0 (0-0.1)
Chile	Intra-abdominal infections	2,640 (1,770-3,790)	11.1 (7.4-16)	58,800 (38,100-84,000)	253.4 (163.4-364.3)	718 (468-1,030)	3 (2-4.4)	16,000 (10,100-23,400)	69.1 (43.1-101.2)
Chile	LRI and thorax infections	2,840 (2,230-3,660)	12.4 (9.7-16)	47,500 (35,900-64,300)	228.3 (169.8-310.3)	756 (546-1,020)	3.3 (2.4-4.4)	12,500 (8,850-17,500)	59.6 (42-83.1)
Chile	TB	33 (21-54)	0.1 (0.1-0.2)	814 (537-1,360)	3.6 (2.4-6)	12 (3-30)	0.1 (0-0.1)	276 (82-671)	1.2 (0.4-3)
Chile	Typhoid, paratyphoid, and iNTS	1 (0-2)	0 (0-0)	15 (2-51)	0.1 (0-0.2)	0 (0-0)	0 (0-0)	3 (0-11)	0 (0-0)

Chile	UTI	1,260 (995-1,570)	5.3 (4.2-6.6)	19,000 (14,800-24,700)	81.2 (63.4-105.8)	318 (244-413)	1.3 (1-1.7)	4,790 (3,650-6,460)	20.5 (15.5-27.8)
Colombia	All	18,200 (11,800-26,500)	34.6 (22.4-50.6)	475,000 (301,000-711,000)	999.8 (634.5-1494.8)	4,720 (3,020-7,060)	9 (5.8-13.5)	123,000 (76,500-186,000)	258 (161.2-390.4)
Colombia	Bacterial skin infections	710 (371-1,320)	1.3 (0.7-2.4)	14,900 (7,790-27,800)	29 (15.3-54.5)	167 (80-317)	0.3 (0.1-0.6)	3,510 (1,720-6,580)	6.8 (3.4-12.8)
Colombia	Bone and joint infections	53 (16-122)	0.1 (0-0.2)	1,340 (399-3,170)	2.7 (0.8-6.3)	14 (4-32)	0 (0-0.1)	344 (100-822)	0.7 (0.2-1.6)
Colombia	BSI	5,890 (3,020-9,990)	11.5 (6-19.3)	179,000 (99,100-284,000)	388.5 (219.1-611.2)	1,550 (775-2,640)	3 (1.6-5.1)	46,700 (25,800-77,400)	101.6 (56.8-166)
Colombia	Cardiac infections	386 (235-556)	0.7 (0.4-1.1)	10,400 (6,460-14,900)	20.5 (12.9-29.4)	101 (60-147)	0.2 (0.1-0.3)	2,710 (1,650-3,910)	5.4 (3.3-7.8)
Colombia	CNS infections	144 (94-223)	0.3 (0.2-0.5)	7,090 (4,510-10,800)	15.9 (10-24)	35 (22-55)	0.1 (0-0.1)	1,690 (1,060-2,650)	3.8 (2.4-5.9)
Colombia	Diarrhoea	22 (12-37)	0 (0-0.1)	1,460 (781-2,410)	3.4 (1.8-5.7)	5 (2-9)	0 (0-0)	269 (113-489)	0.6 (0.3-1.2)
Colombia	Gonorrhoea and chlamydia			99 (38-219)	0.2 (0.1-0.4)			10 (2-24)	0 (0-0)
Colombia	Intra-abdominal infections	3,780 (2,420-5,660)	7.1 (4.5-10.6)	83,200 (50,300-129,000)	162.4 (98.3-254.3)	970 (603-1,490)	1.8 (1.1-2.8)	21,400 (12,700-33,900)	41.8 (24.6-66.6)
Colombia	LRI and thorax infections	5,210 (3,480-7,560)	10 (6.7-14.4)	141,000 (94,100-203,000)	305 (202.9-440.4)	1,380 (874-2,030)	2.6 (1.7-3.9)	36,300 (23,500-54,700)	78.3 (50.6-117.4)
Colombia	TB	121 (44-309)	0.2 (0.1-0.6)	3,900 (1,430-9,870)	7.7 (2.8-19.5)	59 (8-190)	0.1 (0-0.4)	1,700 (242-5,340)	3.4 (0.5-10.5)
Colombia	Typhoid, paratyphoid, and iNTS	2 (1-5)	0 (0-0)	70 (20-179)	0.1 (0-0.4)	0 (0-1)	0 (0-0)	14 (2-41)	0 (0-0.1)
Colombia	UTI	1,840 (1,320-2,540)	3.3 (2.4-4.6)	33,300 (23,200-47,700)	64.3 (44.8-92.2)	446 (306-629)	0.8 (0.6-1.1)	8,070 (5,430-11,700)	15.6 (10.5-22.7)
Costa Rica	All	1,890 (1,210-2,800)	37.3 (23.8-55.3)	44,000 (27,100-67,400)	908.9 (557.9-1403.7)	484 (304-731)	9.5 (6-14.4)	11,200 (6,820-17,500)	230.8 (140.7-360.5)
Costa Rica	Bacterial skin infections	61 (28-122)	1.2 (0.5-2.4)	1,290 (587-2,580)	25.5 (11.8-51)	14 (6-27)	0.3 (0.1-0.5)	285 (125-587)	5.6 (2.5-11.6)
Costa Rica	Bone and joint infections	7 (2-17)	0.1 (0-0.3)	163 (45-394)	3.2 (0.9-7.7)	2 (0-4)	0 (0-0.1)	42 (11-105)	0.8 (0.2-2.1)
Costa Rica	BSI	617 (311-1,040)	12.3 (6.2-20.6)	15,500 (7,940-25,500)	324.8 (169.5-536.5)	160 (81-280)	3.2 (1.6-5.5)	3,990 (2,040-6,900)	83.7 (43.6-144)

Costa Rica	Cardiac infections	52 (31-77)	1 (0.6-1.5)	1,330 (792-1,980)	26.1 (15.6-38.6)	14 (8-21)	0.3 (0.2-0.4)	344 (205-522)	6.7 (4-10.3)
Costa Rica	CNS infections	15 (10-23)	0.3 (0.2-0.5)	622 (410-967)	13.9 (9-21.6)	3 (2-5)	0.1 (0-0.1)	139 (87-219)	3.1 (1.9-4.9)
Costa Rica	Diarrhoea	2 (1-3)	0 (0-0.1)	77 (40-135)	1.8 (0.9-3.2)	0 (0-1)	0 (0-0)	13 (6-25)	0.3 (0.1-0.6)
Costa Rica	Gonorrhoea and chlamydia			17 (7-34)	0.3 (0.1-0.7)			2 (0-4)	0 (0-0.1)
Costa Rica	Intra-abdominal infections	512 (315-772)	10 (6.2-15.1)	11,600 (6,940-17,900)	226.6 (135.4-353.2)	128 (79-198)	2.5 (1.5-3.9)	2,930 (1,720-4,660)	57.1 (33.4-91.1)
Costa Rica	LRI and thorax infections	481 (322-700)	9.5 (6.3-13.9)	10,800 (7,050-15,800)	233.8 (151.4-345.2)	129 (79-195)	2.5 (1.6-3.9)	2,820 (1,740-4,320)	60.7 (37-92.6)
Costa Rica	TB	3 (1-10)	0.1 (0-0.2)	106 (40-289)	2.1 (0.8-5.7)	2 (0-6)	0 (0-0.1)	44 (6-155)	0.8 (0.1-3)
Costa Rica	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Costa Rica	UTI	141 (96-199)	2.7 (1.9-3.8)	2,570 (1,730-3,720)	50.7 (34-73.3)	33 (22-47)	0.6 (0.4-0.9)	600 (390-881)	11.8 (7.7-17.3)
Cuba	All	6,830 (4,700-10,000)	35.9 (24.6-52.9)	135,000 (89,700-202,000)	837.9 (555.4-1245.4)	1,680 (1,120-2,520)	8.8 (5.8-13.3)	33,200 (21,600-50,800)	204.9 (132.9-312.3)
Cuba	Bacterial skin infections	270 (119-542)	1.4 (0.6-2.7)	5,020 (2,430-9,860)	30.2 (15.3-56.8)	55 (22-118)	0.3 (0.1-0.6)	1,020 (441-2,040)	6.1 (2.9-11.8)
Cuba	Bone and joint infections	30 (9-73)	0.2 (0-0.4)	531 (151-1,300)	3 (0.8-7.3)	7 (2-17)	0 (0-0.1)	119 (32-307)	0.7 (0.2-1.7)
Cuba	BSI	2,210 (1,160-3,950)	12 (6.4-21.1)	46,900 (25,100-82,000)	305.6 (166.8-524)	585 (297-1,050)	3.2 (1.6-5.6)	12,300 (6,410-21,500)	79.9 (42.6-136.7)
Cuba	Cardiac infections	86 (58-122)	0.5 (0.3-0.7)	2,140 (1,460-3,070)	13.6 (9.3-19.5)	23 (15-33)	0.1 (0.1-0.2)	559 (376-810)	3.6 (2.4-5.1)
Cuba	CNS infections	45 (30-72)	0.3 (0.2-0.5)	1,620 (1,100-2,490)	15.2 (10.4-22.1)	10 (6-16)	0.1 (0-0.1)	350 (229-549)	3.3 (2.2-5)
Cuba	Diarrhoea	2 (1-4)	0 (0-0)	179 (100-302)	1.4 (0.8-2.4)	1 (0-1)	0 (0-0)	32 (16-57)	0.3 (0.1-0.5)
Cuba	Gonorrhoea and chlamydia			74 (36-146)	0.7 (0.3-1.4)			7 (2-16)	0.1 (0-0.2)
Cuba	Intra-abdominal infections	1,340 (865-2,010)	7.1 (4.6-10.7)	29,700 (18,800-45,200)	171.6 (107.2-263.6)	334 (210-515)	1.8 (1.1-2.7)	7,410 (4,570-11,600)	42.8 (26-67.4)

Cuba	LRI and thorax infections	2,480 (1,900-3,250)	12.6 (9.6-16.7)	41,500 (30,800-56,600)	255.3 (189.5-347.1)	573 (417-785)	2.9 (2.1-4)	9,580 (6,770-13,500)	58.3 (41-81.3)
Cuba	TB	4 (1-8)	0 (0-0)	116 (51-264)	0.8 (0.3-1.7)	2 (0-5)	0 (0-0)	38 (7-116)	0.2 (0-0.7)
Cuba	Typhoid, paratyphoid, and iNTS	0 (0-1)	0 (0-0)	5 (1-16)	0 (0-0.1)	0 (0-0)	0 (0-0)	1 (0-3)	0 (0-0)
Cuba	UTI	369 (229-570)	1.9 (1.2-2.9)	7,060 (4,420-10,800)	40.5 (25.3-61.5)	90 (55-143)	0.5 (0.3-0.7)	1,720 (1,060-2,690)	9.8 (6-15.2)
Dominica	All	60 (39-86)	72 (46.5-103.5)	1,420 (894-2,100)	2061.8 (1299.6-3076.8)	16 (10-23)	18.4 (11.7-27.1)	359 (224-533)	517.9 (324.6-777.9)
Dominica	Bacterial skin infections	5 (3-8)	5.7 (3.5-9.3)	91 (56-147)	109.8 (68.6-173.9)	1 (1-2)	1.3 (0.8-2.2)	21 (12-36)	26 (15.4-42.9)
Dominica	Bone and joint infections	0 (0-1)	0.2 (0.1-0.6)	4 (1-10)	5 (1.3-12.5)	0 (0-0)	0.1 (0-0.2)	1 (0-3)	1.3 (0.3-3.4)
Dominica	BSI	21 (11-36)	26.3 (14.1-43)	550 (304-870)	852.8 (493-1318.8)	6 (3-9)	6.9 (3.7-11.2)	143 (79-230)	220.4 (127.1-346.5)
Dominica	Cardiac infections	1 (1-2)	1.3 (1-1.8)	27 (19-37)	32.9 (23.7-45.3)	0 (0-0)	0.3 (0.2-0.5)	7 (5-10)	8.6 (6.1-11.9)
Dominica	CNS infections	0 (0-1)	0.7 (0.4-1.3)	24 (14-42)	43.1 (24.6-73.9)	0 (0-0)	0.2 (0.1-0.3)	5 (3-10)	9.8 (5.5-17.1)
Dominica	Diarrhoea	0 (0-0)	0 (0-0)	1 (0-2)	1.7 (0.9-3.1)	0 (0-0)	0 (0-0)	0 (0-0)	0.4 (0.2-0.7)
Dominica	Gonorrhoea and chlamydia			1 (0-2)	1.1 (0.4-2.6)			0 (0-0)	0.1 (0-0.3)
Dominica	Intra-abdominal infections	9 (6-15)	10.7 (6.6-16.9)	209 (124-336)	251.4 (146.3-410.5)	2 (1-4)	2.7 (1.6-4.4)	53 (30-88)	64.1 (36.2-106.4)
Dominica	LRI and thorax infections	19 (12-27)	22.7 (15.2-32.9)	442 (286-648)	681.6 (438.3-997.9)	5 (3-7)	5.8 (3.7-8.7)	110 (69-165)	166.3 (104.1-247.2)
Dominica	TB	0 (0-0)	0.1 (0-0.3)	2 (0-8)	2.5 (0.7-9.9)	0 (0-0)	0 (0-0.2)	1 (0-4)	1.2 (0.1-5.8)
Dominica	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-1)	0.4 (0.1-0.9)	0 (0-0)	0 (0-0)	0 (0-0)	0.1 (0-0.2)
Dominica	UTI	4 (2-6)	4.1 (2.3-6.8)	68 (38-112)	79.4 (44.6-131.5)	1 (1-2)	1 (0.6-1.7)	17 (9-28)	19.5 (10.7-32.8)
Dominican Republic	All	5,500 (3,440-8,270)	59.4 (37.4-89.4)	209,000 (133,000-311,000)	2032.6 (1293.7-3032.2)	1,390 (870-2,090)	15 (9.4-22.6)	51,900 (32,700-78,000)	505.2 (317.4-757.6)

Dominican Republic	Bacterial skin infections	174 (49-398)	2 (0.6-4.5)	4,010 (1,270-8,990)	41.8 (13-93.3)	40 (11-92)	0.5 (0.1-1)	924 (278-2,140)	9.6 (2.8-21.8)
Dominican Republic	Bone and joint infections	20 (6-47)	0.2 (0.1-0.5)	442 (116-1,100)	4.6 (1.2-11.4)	5 (1-13)	0.1 (0-0.1)	113 (30-296)	1.2 (0.3-3.1)
Dominican Republic	BSI	1,990 (1,150-3,110)	21.1 (12-33.5)	86,800 (54,900-127,000)	832.8 (530.2-1226.7)	501 (290-796)	5.3 (3-8.6)	21,700 (13,700-32,600)	208.2 (131-313)
Dominican Republic	Cardiac infections	58 (40-81)	0.6 (0.4-0.9)	1,950 (1,300-2,770)	18.7 (12.5-26.5)	15 (10-20)	0.2 (0.1-0.2)	489 (332-698)	4.7 (3.2-6.7)
Dominican Republic	CNS infections	96 (62-148)	0.9 (0.6-1.5)	5,620 (3,450-8,700)	52.1 (32.1-80.5)	22 (14-35)	0.2 (0.1-0.3)	1,310 (792-2,080)	12.1 (7.4-19.3)
Dominican Republic	Diarrhoea	6 (3-12)	0.1 (0-0.1)	525 (223-1,050)	4.9 (2.1-9.6)	2 (1-3)	0 (0-0)	115 (45-242)	1.1 (0.4-2.2)
Dominican Republic	Gonorrhoea and chlamydia			136 (43-402)	1.2 (0.4-3.3)			14 (2-43)	0.1 (0-0.4)
Dominican Republic	Intra-abdominal infections	1,170 (697-1,880)	12.8 (7.7-20.4)	32,700 (18,500-53,600)	330.2 (188.4-537.8)	297 (170-476)	3.2 (1.9-5.1)	8,330 (4,630-13,700)	84 (47.2-137)
Dominican Republic	LRI and thorax infections	1,850 (1,190-2,700)	20.1 (13-29.4)	72,800 (45,300-109,000)	704.3 (442.8-1052.9)	467 (296-688)	5.1 (3.3-7.5)	17,800 (10,800-27,200)	173.2 (105.8-262.4)
Dominican Republic	TB	29 (14-70)	0.3 (0.1-0.7)	1,280 (634-3,030)	11.7 (5.8-27.7)	10 (2-35)	0.1 (0-0.3)	416 (79-1,440)	3.8 (0.7-13.2)
Dominican Republic	Typhoid, paratyphoid, and iNTS	2 (1-4)	0 (0-0)	82 (28-192)	0.8 (0.3-1.8)	0 (0-1)	0 (0-0)	16 (3-44)	0.1 (0-0.4)
Dominican Republic	UTI	114 (68-185)	1.3 (0.8-2.1)	2,830 (1,710-4,480)	29.3 (17.8-46.6)	28 (16-46)	0.3 (0.2-0.5)	675 (397-1,110)	7 (4.1-11.6)
Ecuador	All	9,290 (6,430-13,100)	66.9 (46.4-94.1)	272,000 (187,000-382,000)	1716.6 (1186.6-2388.9)	2,410 (1,620-3,440)	17.3 (11.7-24.6)	69,800 (47,200-99,700)	440.2 (298-625.8)
Ecuador	Bacterial skin infections	187 (65-406)	1.4 (0.5-2.9)	4,160 (1,650-8,890)	27.2 (10.7-58)	42 (14-94)	0.3 (0.1-0.7)	921 (344-1,980)	6 (2.3-12.9)
Ecuador	Bone and joint infections	27 (7-62)	0.2 (0.1-0.4)	633 (174-1,520)	4.1 (1.1-9.6)	7 (2-16)	0 (0-0.1)	156 (42-383)	1 (0.3-2.5)

Ecuador	BSI	2,670 (1,460-4,370)	18.7 (10-31.1)	87,200 (52,700-135,000)	543.2 (325.5-845.9)	695 (380-1,130)	4.9 (2.6-8)	22,500 (13,500-35,400)	140.5 (83.2-220.5)
Ecuador	Cardiac infections	65 (48-89)	0.4 (0.3-0.6)	2,100 (1,540-2,920)	12.7 (9.3-17.4)	17 (12-24)	0.1 (0.1-0.2)	542 (384-762)	3.3 (2.3-4.6)
Ecuador	CNS infections	67 (42-116)	0.4 (0.3-0.7)	3,460 (2,170-5,830)	20.1 (12.7-34.1)	16 (10-29)	0.1 (0.1-0.2)	841 (512-1,430)	4.9 (3-8.4)
Ecuador	Diarrhoea	3 (1-6)	0 (0-0)	259 (118-518)	1.6 (0.7-3.1)	1 (0-1)	0 (0-0)	50 (24-96)	0.3 (0.1-0.6)
Ecuador	Gonorrhoea and chlamydia			74 (38-134)	0.4 (0.2-0.7)			7 (2-16)	0 (0-0.1)
Ecuador	Intra-abdominal infections	2,180 (1,400-3,180)	15.3 (9.9-22.3)	51,400 (31,500-77,600)	330.7 (204.1-495.3)	564 (349-849)	4 (2.5-6)	13,300 (7,940-20,500)	85.7 (51.4-131.8)
Ecuador	LRI and thorax infections	3,490 (2,640-4,590)	25.9 (19.7-33.9)	108,000 (79,600-144,000)	680.9 (502.9-903.9)	886 (637-1,210)	6.6 (4.8-9)	26,900 (19,000-36,400)	169.8 (120.8-229)
Ecuador	TB	128 (48-322)	0.8 (0.3-2.1)	4,820 (1,850-11,900)	28.4 (10.9-70.3)	59 (9-188)	0.4 (0.1-1.2)	2,050 (317-6,340)	12.1 (1.9-37.6)
Ecuador	Typhoid, paratyphoid, and iNTS	1 (0-2)	0 (0-0)	54 (19-114)	0.3 (0.1-0.6)	0 (0-0)	0 (0-0)	11 (2-28)	0.1 (0-0.2)
Ecuador	UTI	479 (305-752)	3.7 (2.3-5.7)	10,100 (6,360-16,000)	67 (42.9-105.3)	120 (75-193)	0.9 (0.6-1.5)	2,480 (1,530-4,070)	16.5 (10.2-26.9)
El Salvador	All	3,390 (2,210-4,930)	54.5 (35.3-79.6)	85,400 (53,600-129,000)	1439.8 (900.2-2177)	857 (546-1,290)	13.8 (8.7-20.8)	21,400 (13,200-33,100)	360.5 (221.7-558.1)
El Salvador	Bacterial skin infections	100 (45-204)	1.6 (0.7-3.3)	2,120 (933-4,330)	35.2 (15.3-72.3)	23 (10-48)	0.4 (0.2-0.8)	484 (203-1,020)	8 (3.3-17.1)
El Salvador	Bone and joint infections	12 (3-29)	0.2 (0.1-0.5)	266 (66-668)	4.4 (1.1-11.2)	3 (1-8)	0 (0-0.1)	66 (17-175)	1.1 (0.3-2.9)
El Salvador	BSI	995 (496-1,650)	16.3 (8.2-27)	27,800 (15,000-45,400)	474 (258-770.7)	253 (129-433)	4.1 (2.1-7)	7,030 (3,800-11,800)	119.7 (65.1-199.6)
El Salvador	Cardiac infections	18 (11-28)	0.3 (0.2-0.5)	543 (341-856)	9 (5.7-14.2)	5 (3-7)	0.1 (0-0.1)	137 (86-212)	2.3 (1.4-3.5)
El Salvador	CNS infections	21 (14-34)	0.4 (0.2-0.6)	872 (549-1,380)	14.5 (9.1-23.2)	5 (3-8)	0.1 (0.1-0.1)	202 (124-321)	3.4 (2-5.4)
El Salvador	Diarrhoea	4 (2-8)	0.1 (0-0.1)	230 (105-454)	3.9 (1.8-7.8)	1 (0-2)	0 (0-0)	49 (22-94)	0.8 (0.4-1.6)
El Salvador	Gonorrhoea and chlamydia			15 (6-28)	0.2 (0.1-0.4)			1 (0-3)	0 (0-0.1)

El Salvador	Intra-abdominal infections	686 (404-1,080)	11.3 (6.7-17.7)	17,000 (9,490-27,300)	285.7 (159.8-459.1)	172 (100-274)	2.8 (1.6-4.5)	4,280 (2,380-7,030)	72 (39.9-118)
El Salvador	LRI and thorax infections	1,280 (914-1,730)	20.1 (14.2-27.5)	30,700 (20,800-43,600)	516.9 (347.8-737.4)	328 (222-462)	5.1 (3.5-7.3)	7,700 (5,040-11,200)	129.4 (84.2-188.2)
El Salvador	TB	7 (3-19)	0.1 (0-0.3)	245 (91-627)	4 (1.5-10.3)	3 (0-11)	0.1 (0-0.2)	90 (12-316)	1.5 (0.2-5.2)
El Salvador	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	7 (1-19)	0.1 (0-0.3)	0 (0-0)	0 (0-0)	1 (0-4)	0 (0-0.1)
El Salvador	UTI	269 (179-395)	4.3 (2.8-6.3)	5,520 (3,590-8,270)	91.8 (59.2-137.8)	65 (42-98)	1 (0.7-1.6)	1,340 (848-2,040)	22.3 (14-34)
Grenada	All	72 (50-99)	74.2 (52.4-101.2)	1,860 (1,280-2,600)	1859.7 (1281.3-2575.4)	18 (12-26)	18.8 (13-26.3)	468 (317-669)	465.9 (316.7-662.4)
Grenada	Bacterial skin infections	6 (3-9)	6 (3.6-9.8)	122 (74-198)	117.4 (72.4-190.8)	1 (1-2)	1.3 (0.8-2.3)	28 (16-45)	26.6 (15.3-44)
Grenada	Bone and joint infections	0 (0-1)	0.3 (0.1-0.7)	7 (2-16)	6.2 (1.7-14.8)	0 (0-0)	0.1 (0-0.2)	2 (0-4)	1.6 (0.4-3.8)
Grenada	BSI	22 (12-36)	22.5 (12.3-37.2)	601 (342-951)	612.3 (355.2-946.2)	6 (3-10)	5.9 (3.2-9.8)	156 (87-249)	158.4 (90.3-250.7)
Grenada	Cardiac infections	2 (1-2)	1.7 (1.3-2.4)	47 (36-68)	43 (33.1-60.7)	0 (0-1)	0.4 (0.3-0.6)	12 (9-18)	11.2 (8.4-16)
Grenada	CNS infections	0 (0-1)	0.3 (0.2-0.6)	15 (9-28)	15.5 (9-28.3)	0 (0-0)	0.1 (0-0.2)	3 (2-6)	3.6 (2-6.7)
Grenada	Diarrhoea	0 (0-0)	0 (0-0)	1 (1-2)	1.3 (0.7-2.2)	0 (0-0)	0 (0-0)	0 (0-0)	0.3 (0.1-0.5)
Grenada	Gonorrhoea and chlamydia			1 (1-2)	0.9 (0.5-1.7)			0 (0-0)	0.1 (0-0.2)
Grenada	Intra-abdominal infections	13 (8-19)	12 (7.8-18)	322 (202-498)	287.1 (180.7-441.9)	3 (2-5)	3.1 (1.9-4.7)	83 (50-129)	73.5 (45-114.8)
Grenada	LRI and thorax infections	23 (17-31)	25.2 (19-33.4)	614 (451-827)	654.5 (474.4-879.8)	6 (4-8)	6.3 (4.6-8.6)	152 (107-210)	160.8 (112.8-222)
Grenada	TB	0 (0-0)	0 (0-0.1)	1 (0-2)	0.6 (0.2-2.2)	0 (0-0)	0 (0-0)	0 (0-1)	0.2 (0-0.9)
Grenada	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0.1)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Grenada	UTI	6 (4-9)	6.2 (4.5-8.8)	129 (91-184)	121 (86-172.2)	1 (1-2)	1.5 (1.1-2.2)	32 (22-47)	29.6 (20.4-43.6)

Guatemala	All	11,800 (8,090-16,500)	104.5 (72.9-145.7)	438,000 (298,000-628,000)	2919.2 (1991.2-4164.1)	2,850 (1,920-4,030)	25.5 (17.4-35.6)	104,000 (68,900-150,000)	699.2 (465.7-996.1)
Guatemala	Bacterial skin infections	272 (110-564)	2.7 (1.1-5.5)	7,140 (3,080-14,000)	56.7 (24.1-112.4)	62 (24-129)	0.6 (0.2-1.3)	1,630 (680-3,310)	13 (5.2-26.9)
Guatemala	Bone and joint infections	32 (9-76)	0.3 (0.1-0.7)	830 (218-2,090)	6.5 (1.7-16)	8 (2-18)	0.1 (0-0.2)	201 (53-515)	1.6 (0.4-4)
Guatemala	BSI	2,630 (1,420-4,270)	23 (11.8-37.7)	101,000 (58,500-160,000)	666.6 (374.1-1059.4)	654 (357-1,080)	5.7 (3-9.7)	24,800 (14,100-39,600)	164.5 (92.5-264.5)
Guatemala	Cardiac infections	42 (30-59)	0.3 (0.2-0.5)	1,800 (1,240-2,530)	11.3 (8-15.5)	10 (7-15)	0.1 (0.1-0.1)	438 (296-615)	2.8 (1.9-3.8)
Guatemala	CNS infections	101 (70-149)	0.7 (0.5-1)	5,360 (3,600-8,030)	31.1 (20.9-46.1)	23 (15-34)	0.2 (0.1-0.2)	1,210 (773-1,840)	7 (4.5-10.6)
Guatemala	Diarrhoea	38 (20-64)	0.3 (0.2-0.5)	2,330 (1,230-4,040)	13 (7.1-22.5)	9 (4-17)	0.1 (0-0.1)	543 (251-1,040)	3 (1.4-5.7)
Guatemala	Gonorrhoea and chlamydia			75 (30-162)	0.4 (0.2-0.8)			7 (2-18)	0 (0-0.1)
Guatemala	Intra-abdominal infections	2,400 (1,450-3,620)	21.2 (13-31.5)	74,900 (43,100-118,000)	563.6 (327.4-871.7)	608 (362-957)	5.4 (3.2-8.4)	19,000 (10,900-30,400)	142.7 (83.1-228.9)
Guatemala	LRI and thorax infections	5,720 (4,280-7,490)	51 (38.9-65.5)	230,000 (167,000-313,000)	1457.3 (1066.9-1951.6)	1,340 (957-1,820)	12.1 (8.8-16.1)	52,400 (36,700-73,600)	335 (236.6-462.5)
Guatemala	TB	50 (22-109)	0.4 (0.2-0.8)	2,190 (969-4,780)	14.2 (6.2-30.8)	20 (3-59)	0.1 (0-0.4)	816 (128-2,510)	5.3 (0.8-15.9)
Guatemala	Typhoid, paratyphoid, and iNTS	17 (3-33)	0.1 (0-0.2)	851 (169-1,770)	4.9 (1-9.9)	3 (0-8)	0 (0-0.1)	167 (15-428)	1 (0.1-2.5)
Guatemala	UTI	457 (276-716)	4.5 (2.8-7.2)	11,600 (7,150-18,100)	93.5 (56.9-146.7)	114 (67-182)	1.1 (0.7-1.8)	2,890 (1,730-4,590)	23.4 (13.7-37.3)
Guyana	All	579 (373-855)	102.7 (67.6-148.4)	19,300 (12,200-29,100)	2895.5 (1836.6-4327.8)	147 (93-218)	26.2 (16.9-38)	4,860 (3,030-7,290)	729 (458.4-1084.4)
Guyana	Bacterial skin infections	21 (7-43)	3.9 (1.4-8.1)	519 (185-1,100)	81.6 (28.7-169.2)	5 (2-11)	0.9 (0.3-2)	122 (41-268)	19 (6.4-42.1)
Guyana	Bone and joint infections	2 (0-5)	0.4 (0.1-0.9)	51 (12-129)	7.8 (1.9-19.7)	0 (0-1)	0.1 (0-0.2)	13 (3-34)	2 (0.5-5.2)
Guyana	BSI	155 (82-254)	26.9 (14.1-44.5)	5,620 (3,220-8,930)	839.3 (479.7-1341.9)	39 (21-65)	6.7 (3.5-11.4)	1,390 (798-2,230)	208.4 (119.4-333.5)
Guyana	Cardiac infections	11 (8-14)	1.7 (1.2-2.3)	369 (255-506)	51.2 (36-69.6)	3 (2-4)	0.4 (0.3-0.6)	91 (63-128)	12.7 (8.9-17.6)

Guyana	CNS infections	4 (2-8)	0.6 (0.3-1.2)	219 (120-397)	29.5 (16.3-53.6)	1 (1-2)	0.1 (0.1-0.3)	51 (27-94)	6.9 (3.7-12.8)
Guyana	Diarrhoea	0 (0-1)	0.1 (0-0.1)	32 (17-52)	4.6 (2.5-7.6)	0 (0-0)	0 (0-0)	7 (4-12)	1.1 (0.6-1.8)
Guyana	Gonorrhoea and chlamydia			8 (3-16)	0.9 (0.4-1.7)			1 (0-2)	0.1 (0-0.2)
Guyana	Intra-abdominal infections	116 (69-178)	18.9 (11.5-28.4)	3,540 (2,030-5,480)	506.8 (296.6-785.3)	30 (18-47)	4.8 (2.9-7.6)	907 (508-1,470)	129.8 (73.8-208.8)
Guyana	LRI and thorax infections	240 (163-344)	44.8 (30.8-63)	8,220 (5,340-11,900)	1255.2 (827.4-1813.3)	62 (40-90)	11.7 (7.6-16.6)	2,080 (1,310-3,050)	318.7 (200.4-466)
Guyana	TB	2 (1-6)	0.2 (0.1-0.9)	66 (22-262)	8.7 (2.9-34.7)	1 (0-4)	0.1 (0-0.5)	28 (3-142)	3.8 (0.4-18.9)
Guyana	Typhoid, paratyphoid, and iNTS	0 (0-1)	0 (0-0.1)	11 (3-32)	1.5 (0.4-4.2)	0 (0-0)	0 (0-0)	2 (0-7)	0.3 (0-0.9)
Guyana	UTI	27 (19-39)	5.2 (3.6-7.3)	691 (468-989)	108.4 (73.8-154.3)	7 (4-10)	1.3 (0.8-1.8)	168 (111-246)	26.3 (17.5-38.3)
Haiti	All	10,200 (7,220-14,000)	121.3 (84-170.6)	564,000 (413,000-751,000)	4574.1 (3335.7-6210.7)	2,470 (1,700-3,460)	30 (20-42.4)	135,000 (95,600-180,000)	1102.1 (771-1514.7)
Haiti	Bacterial skin infections	382 (181-717)	6.6 (3-12.4)	11,300 (5,640-21,000)	138.2 (67.6-255.4)	84 (38-160)	1.4 (0.6-2.8)	2,490 (1,200-4,680)	30.2 (14.1-56.7)
Haiti	Bone and joint infections	28 (7-71)	0.4 (0.1-1.1)	1,050 (259-2,820)	10.9 (2.8-27.8)	7 (2-17)	0.1 (0-0.3)	253 (60-681)	2.6 (0.6-6.8)
Haiti	BSI	2,650 (1,630-4,010)	31.3 (17.8-50.3)	145,000 (96,000-211,000)	1178.5 (751.2-1761.1)	664 (411-1,020)	7.9 (4.5-12.7)	36,200 (23,200-53,900)	294.9 (185-442.4)
Haiti	Cardiac infections	59 (39-91)	0.7 (0.5-1.1)	2,590 (1,670-4,120)	24 (16-37)	15 (10-22)	0.2 (0.1-0.3)	641 (410-1,020)	6 (4-9.1)
Haiti	CNS infections	637 (403-965)	4.8 (3-7.6)	50,800 (31,900-77,800)	344.6 (218.3-528.8)	149 (90-238)	1.1 (0.7-1.8)	11,900 (7,140-18,500)	80.3 (48.6-126.5)
Haiti	Diarrhoea	60 (26-120)	0.5 (0.2-0.9)	5,030 (2,060-10,300)	33.8 (14.6-67.7)	15 (6-30)	0.1 (0.1-0.2)	1,210 (470-2,570)	8.2 (3.3-17.1)
Haiti	Gonorrhoea and chlamydia			143 (46-406)	1.1 (0.4-2.9)			14 (3-42)	0.1 (0-0.3)
Haiti	Intra-abdominal infections	1,400 (775-2,210)	20.2 (11.7-31.1)	48,400 (24,500-83,800)	530.7 (287.8-862.1)	352 (196-564)	5.1 (3-8)	12,200 (5,950-21,100)	133.5 (70.8-217.9)
Haiti	LRI and thorax infections	4,660 (3,430-6,180)	51.9 (37.1-71.9)	290,000 (214,000-375,000)	2201.9 (1632.1-2882.4)	1,110 (781-1,500)	12.8 (8.8-17.9)	67,100 (47,600-89,700)	517.8 (363.7-692.2)

Haiti	TB	23 (9-68)	0.3 (0.1-0.8)	1,130 (423-3,420)	10 (3.8-29.5)	10 (2-40)	0.1 (0-0.5)	462 (73-1,870)	4.1 (0.6-16.3)
Haiti	Typhoid, paratyphoid, and iNTS	2 (1-5)	0 (0-0)	153 (43-370)	1 (0.3-2.5)	0 (0-1)	0 (0-0)	30 (4-93)	0.2 (0-0.6)
Haiti	UTI	270 (179-427)	4.6 (3-7.5)	8,310 (5,610-12,200)	99.5 (65.9-152.9)	66 (43-103)	1.1 (0.7-1.8)	2,020 (1,360-2,940)	24.2 (15.9-36.8)
Honduras	All	5,240 (3,530-7,380)	90.4 (61.4-126.2)	170,000 (111,000-246,000)	2235.5 (1492.7-3204.2)	1,320 (886-1,880)	22.8 (15.5-32.4)	41,900 (27,500-61,100)	556.4 (368.6-806.9)
Honduras	Bacterial skin infections	125 (56-256)	2.4 (1.1-4.8)	3,080 (1,440-5,990)	47.6 (22.2-93.9)	29 (12-58)	0.6 (0.2-1.1)	703 (314-1,440)	10.9 (4.8-21.9)
Honduras	Bone and joint infections	16 (5-35)	0.3 (0.1-0.7)	392 (110-916)	6 (1.7-13.5)	4 (1-9)	0.1 (0-0.2)	97 (28-239)	1.5 (0.4-3.5)
Honduras	BSI	1,500 (876-2,340)	25.2 (14-40.8)	52,000 (32,000-76,900)	661.7 (397.6-991)	371 (210-594)	6.2 (3.4-10.3)	12,700 (7,830-19,400)	162.4 (97.8-250.6)
Honduras	Cardiac infections	32 (21-45)	0.6 (0.4-0.8)	840 (532-1,270)	12.4 (8.1-18)	8 (5-11)	0.1 (0.1-0.2)	205 (127-314)	3 (1.9-4.5)
Honduras	CNS infections	48 (31-75)	0.7 (0.5-1.1)	2,300 (1,420-3,700)	25.6 (16.4-40.9)	11 (7-17)	0.2 (0.1-0.2)	507 (302-828)	5.7 (3.5-9.2)
Honduras	Diarrhoea	22 (10-42)	0.3 (0.1-0.6)	1,560 (741-3,020)	15.3 (7.2-29)	5 (2-10)	0.1 (0-0.1)	347 (138-704)	3.4 (1.4-6.7)
Honduras	Gonorrhoea and chlamydia			38 (14-90)	0.4 (0.1-0.9)			4 (1-11)	0 (0-0.1)
Honduras	Intra-abdominal infections	1,620 (1,040-2,420)	28.2 (18.1-41.3)	46,000 (28,500-69,000)	656.4 (411-989.9)	407 (254-597)	7.1 (4.4-10.4)	11,600 (6,960-17,600)	165 (101.4-247)
Honduras	LRI and thorax infections	1,480 (959-2,140)	25.5 (16.5-37)	53,700 (34,400-77,900)	659.9 (431.5-951.3)	374 (234-554)	6.5 (4-9.8)	13,000 (8,150-19,200)	163.2 (103.1-241.3)
Honduras	TB	77 (37-158)	1.3 (0.6-2.6)	2,530 (1,190-5,160)	34.6 (16.7-69.9)	30 (7-87)	0.5 (0.1-1.4)	935 (208-2,690)	12.9 (2.9-37.1)
Honduras	Typhoid, paratyphoid, and iNTS	0 (0-1)	0 (0-0)	11 (2-30)	0.1 (0-0.3)	0 (0-0)	0 (0-0)	2 (0-7)	0 (0-0.1)
Honduras	UTI	316 (185-494)	6.1 (3.5-9.4)	7,240 (4,130-11,700)	115.5 (67.5-183.3)	78 (44-124)	1.5 (0.8-2.3)	1,780 (996-2,890)	28.4 (15.9-45.7)
Jamaica	All	1,430 (907-2,110)	47.5 (29.9-70.4)	34,900 (21,300-53,300)	1342.7 (816.7-2060)	376 (232-566)	12.5 (7.7-18.7)	9,110 (5,520-13,900)	349.4 (213.7-543.5)
Jamaica	Bacterial skin infections	117 (61-203)	3.6 (1.8-6.5)	2,100 (1,070-3,780)	70.3 (35.6-127.7)	27 (13-49)	0.8 (0.4-1.5)	486 (239-891)	16.3 (7.9-30)

Jamaica	Bone and joint infections	5 (1-13)	0.2 (0-0.4)	116 (29-288)	3.9 (1-9.7)	1 (0-4)	0 (0-0.1)	31 (8-80)	1 (0.3-2.7)
Jamaica	BSI	551 (280-942)	18.8 (9.8-31.3)	14,500 (8,010-23,200)	577 (328.4-901.1)	150 (76-257)	5.1 (2.7-8.5)	3,910 (2,170-6,310)	154.7 (87.4-243.6)
Jamaica	Cardiac infections	28 (16-38)	0.9 (0.5-1.2)	751 (444-1,030)	25.1 (15-34.6)	8 (4-11)	0.2 (0.1-0.3)	203 (115-288)	6.8 (3.9-9.6)
Jamaica	CNS infections	15 (8-29)	0.5 (0.3-0.9)	643 (363-1,170)	25.4 (14.1-45.7)	4 (2-7)	0.1 (0.1-0.2)	154 (85-285)	6.1 (3.3-11.1)
Jamaica	Diarrhoea	0 (0-1)	0 (0-0)	24 (12-42)	1 (0.5-1.7)	0 (0-0)	0 (0-0)	5 (2-9)	0.2 (0.1-0.4)
Jamaica	Gonorrhoea and chlamydia			23 (9-50)	0.7 (0.3-1.6)			2 (1-6)	0.1 (0-0.2)
Jamaica	Intra-abdominal infections	190 (111-309)	6.2 (3.7-10.2)	4,400 (2,510-7,430)	150.2 (86-253.2)	49 (28-83)	1.6 (0.9-2.7)	1,150 (633-2,010)	39.3 (21.6-68.4)
Jamaica	LRI and thorax infections	389 (239-604)	13.1 (7.9-20.4)	9,970 (5,940-15,800)	411.4 (243.2-648.8)	105 (62-164)	3.5 (2.1-5.6)	2,600 (1,530-4,120)	105.9 (61.4-166)
Jamaica	TB	0 (0-1)	0 (0-0)	15 (3-52)	0.5 (0.1-1.7)	0 (0-1)	0 (0-0)	7 (1-28)	0.2 (0-0.9)
Jamaica	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-1)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Jamaica	UTI	134 (88-186)	4.2 (2.7-5.9)	2,310 (1,560-3,250)	77 (52.2-108.7)	33 (21-47)	1 (0.7-1.5)	566 (378-815)	18.9 (12.6-27.2)
Mexico	All	70,100 (47,400-100,000)	63 (42.9-89.6)	1,970,000 (1,350,000-2,780,000)	1703.9 (1178.7-2384.6)	18,300 (12,400-26,100)	16.4 (11.3-23.4)	511,000 (349,000-727,000)	440.6 (302.9-624.6)
Mexico	Bacterial skin infections	2,240 (985-4,420)	2 (0.9-4)	51,500 (23,700-98,900)	43.5 (20.1-84.5)	529 (228-1,050)	0.5 (0.2-0.9)	12,200 (5,510-24,100)	10.3 (4.6-20.2)
Mexico	Bone and joint infections	232 (67-541)	0.2 (0.1-0.5)	5,710 (1,580-13,800)	4.8 (1.3-11.4)	59 (16-141)	0.1 (0-0.1)	1,430 (395-3,540)	1.2 (0.3-3)
Mexico	BSI	21,700 (11,900-35,500)	19.6 (10.8-32)	659,000 (386,000-1,020,000)	579.4 (344.6-886)	5,690 (3,130-9,380)	5.1 (2.8-8.5)	172,000 (101,000-269,000)	150.8 (90.6-234.1)
Mexico	Cardiac infections	425 (307-584)	0.4 (0.3-0.5)	13,600 (9,860-18,600)	11.2 (8.1-15.3)	111 (78-155)	0.1 (0.1-0.1)	3,520 (2,490-4,850)	2.9 (2.1-4)
Mexico	CNS infections	293 (183-521)	0.2 (0.2-0.4)	13,600 (8,640-22,600)	11.4 (7.2-19.1)	72 (44-130)	0.1 (0-0.1)	3,310 (2,020-5,730)	2.8 (1.7-4.8)
Mexico	Diarrhoea	47 (28-78)	0 (0-0.1)	2,670 (1,480-4,460)	2.5 (1.4-4.2)	12 (7-21)	0 (0-0)	662 (348-1,150)	0.6 (0.3-1.1)

Mexico	Gonorrhoea and chlamydia			576 (314-939)	0.4 (0.2-0.7)			57 (16-118)	0 (0-0.1)
Mexico	Intra-abdominal infections	20,500 (13,300-28,800)	17.8 (11.7-25)	533,000 (333,000-774,000)	439 (276.4-630.4)	5,360 (3,450-7,890)	4.7 (3-6.8)	140,000 (87,400-212,000)	115 (72.3-173.3)
Mexico	LRI and thorax infections	17,800 (12,600-24,400)	16.5 (11.7-22.6)	536,000 (397,000-714,000)	478 (353.1-634.2)	4,620 (3,170-6,490)	4.3 (2.9-6)	137,000 (97,300-186,000)	121.8 (86.5-165.7)
Mexico	TB	318 (156-688)	0.3 (0.1-0.6)	11,700 (5,790-25,200)	9.2 (4.6-20)	133 (26-397)	0.1 (0-0.3)	4,480 (899-13,400)	3.5 (0.7-10.6)
Mexico	Typhoid, paratyphoid, and iNTS	26 (4-141)	0 (0-0.1)	972 (164-4,650)	0.8 (0.1-3.9)	5 (0-25)	0 (0-0)	183 (19-806)	0.2 (0-0.7)
Mexico	UTI	6,540 (4,690-9,090)	6 (4.3-8.2)	145,000 (103,000-204,000)	123.7 (87.8-173.4)	1,670 (1,160-2,410)	1.5 (1.1-2.2)	37,200 (25,500-53,400)	31.6 (21.8-45.3)
Nicaragua	All	2,460 (1,640-3,530)	59.1 (39.4-85.1)	80,300 (55,000-115,000)	1548.6 (1055.3-2201.5)	630 (408-917)	15.2 (9.9-22)	20,400 (13,600-29,500)	393.7 (261.2-571.4)
Nicaragua	Bacterial skin infections	80 (33-166)	2.1 (0.9-4.2)	2,010 (925-3,920)	42.6 (19.1-84.6)	18 (7-38)	0.5 (0.2-1)	455 (201-938)	9.6 (4.1-20)
Nicaragua	Bone and joint infections	9 (2-21)	0.2 (0.1-0.5)	224 (66-519)	4.6 (1.4-10.9)	2 (1-5)	0.1 (0-0.1)	57 (15-141)	1.2 (0.3-2.9)
Nicaragua	BSI	895 (498-1,450)	21.2 (11.4-35.1)	30,300 (18,500-46,600)	578.1 (342.8-893.9)	230 (126-376)	5.5 (2.9-9.1)	7,710 (4,630-12,000)	147.7 (86.6-234)
Nicaragua	Cardiac infections	24 (16-35)	0.6 (0.4-0.8)	744 (511-1,080)	14.1 (9.7-20.7)	6 (4-9)	0.1 (0.1-0.2)	189 (126-285)	3.6 (2.4-5.4)
Nicaragua	CNS infections	19 (12-31)	0.3 (0.2-0.6)	1,070 (700-1,710)	17.2 (11.2-27.9)	4 (3-8)	0.1 (0.1-0.1)	251 (158-405)	4 (2.5-6.6)
Nicaragua	Diarrhoea	4 (2-7)	0.1 (0-0.2)	259 (111-529)	4.4 (1.9-8.9)	1 (0-2)	0 (0-0)	60 (25-125)	1 (0.4-2.1)
Nicaragua	Gonorrhoea and chlamydia			23 (11-46)	0.3 (0.2-0.6)			2 (1-5)	0 (0-0.1)
Nicaragua	Intra-abdominal infections	630 (391-923)	14.9 (9.4-21.7)	17,100 (10,200-25,800)	349.9 (212.3-519)	161 (99-244)	3.8 (2.4-5.7)	4,390 (2,580-6,860)	89.6 (54-137.9)
Nicaragua	LRI and thorax infections	647 (475-880)	15.9 (11.5-21.7)	24,900 (18,800-32,700)	457.9 (344.5-606)	168 (116-236)	4.2 (2.8-5.9)	6,260 (4,420-8,410)	116.4 (81.6-157.9)
Nicaragua	TB	12 (5-26)	0.3 (0.1-0.6)	445 (200-976)	8 (3.6-17.5)	5 (1-14)	0.1 (0-0.3)	162 (23-486)	2.9 (0.4-8.8)

Nicaragua	Typhoid, paratyphoid, and iNTS	0 (0-1)	0 (0-0)	10 (2-25)	0.2 (0-0.4)	0 (0-0)	0 (0-0)	2 (0-6)	0 (0-0.1)
Nicaragua	UTI	137 (84-221)	3.6 (2.2-5.7)	3,270 (2,020-5,310)	71.3 (44.1-115.1)	34 (20-55)	0.9 (0.5-1.4)	810 (491-1,330)	17.6 (10.7-28.9)
Panama	All	1,530 (1,000-2,270)	36.6 (23.9-54.4)	43,400 (28,000-65,000)	1076.7 (694-1609.8)	382 (245-572)	9.1 (5.8-13.7)	10,700 (6,800-16,100)	264.9 (169-398.4)
Panama	Bacterial skin infections	62 (30-118)	1.5 (0.7-2.8)	1,260 (601-2,410)	30.4 (14.4-58.1)	14 (6-27)	0.3 (0.1-0.6)	276 (125-551)	6.6 (3-13.2)
Panama	Bone and joint infections	4 (1-11)	0.1 (0-0.3)	102 (27-255)	2.5 (0.6-6.1)	1 (0-3)	0 (0-0.1)	25 (6-64)	0.6 (0.2-1.5)
Panama	BSI	521 (272-876)	12.6 (6.6-21.1)	16,100 (9,110-25,700)	403.3 (230.4-640.8)	131 (68-225)	3.2 (1.7-5.4)	3,990 (2,310-6,480)	99.8 (58-161.9)
Panama	Cardiac infections	26 (16-36)	0.6 (0.4-0.9)	732 (478-1,030)	17.6 (11.5-24.6)	6 (4-9)	0.2 (0.1-0.2)	182 (115-261)	4.4 (2.8-6.3)
Panama	CNS infections	13 (9-20)	0.3 (0.2-0.5)	706 (452-1,070)	17.7 (11.3-26.9)	3 (2-5)	0.1 (0-0.1)	161 (98-247)	4 (2.5-6.2)
Panama	Diarrhoea	3 (2-6)	0.1 (0-0.1)	240 (119-436)	6.1 (3-11.2)	1 (0-1)	0 (0-0)	51 (23-97)	1.3 (0.6-2.5)
Panama	Gonorrhoea and chlamydia			14 (6-30)	0.3 (0.1-0.7)			1 (0-3)	0 (0-0.1)
Panama	Intra-abdominal infections	304 (187-472)	7.3 (4.5-11.3)	7,400 (4,300-11,900)	179 (103.9-287.8)	76 (45-122)	1.8 (1.1-2.9)	1,860 (1,050-3,070)	44.9 (25.3-74.2)
Panama	LRI and thorax infections	461 (322-641)	10.9 (7.6-15.3)	13,700 (9,430-19,400)	345.2 (236.3-488.8)	113 (76-166)	2.7 (1.8-4)	3,280 (2,170-4,710)	82.2 (54.4-117.9)
Panama	TB	21 (9-52)	0.5 (0.2-1.2)	807 (339-2,000)	19.4 (8.2-48.2)	9 (2-32)	0.2 (0-0.8)	318 (68-1,160)	7.7 (1.6-27.9)
Panama	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Panama	UTI	118 (76-174)	2.7 (1.8-4.1)	2,300 (1,470-3,490)	55.1 (35.3-84)	29 (18-44)	0.7 (0.4-1)	557 (346-859)	13.4 (8.3-20.7)
Paraguay	All	3,050 (2,010-4,490)	56.1 (37-81.9)	90,400 (58,100-134,000)	1526.4 (983.7-2261.1)	786 (508-1,170)	14.4 (9.4-21.3)	23,100 (14,400-34,500)	389.7 (244.5-579.4)
Paraguay	Bacterial skin infections	104 (46-211)	2 (0.8-4)	2,420 (1,120-4,690)	41.8 (19-82.6)	23 (10-48)	0.4 (0.2-0.9)	530 (239-1,090)	9.1 (4-19)
Paraguay	Bone and joint infections	11 (3-26)	0.2 (0.1-0.5)	287 (81-685)	4.8 (1.3-11.6)	3 (1-7)	0.1 (0-0.1)	71 (20-182)	1.2 (0.3-3)

Paraguay	BSI	971 (519-1,590)	17.6 (9.4-29.1)	35,000 (20,300-55,000)	585.5 (341.4-919)	250 (136-420)	4.5 (2.4-7.7)	8,950 (5,240-14,500)	149.9 (87.5-241.3)
Paraguay	Cardiac infections	54 (34-77)	1 (0.6-1.4)	1,480 (961-2,110)	24.4 (15.7-34.6)	14 (9-20)	0.3 (0.2-0.4)	380 (244-546)	6.3 (4-9.1)
Paraguay	CNS infections	24 (16-37)	0.4 (0.3-0.6)	1,390 (880-2,080)	21.5 (13.5-32.2)	6 (4-9)	0.1 (0.1-0.1)	325 (202-494)	5 (3.1-7.7)
Paraguay	Diarrhoea	2 (1-3)	0 (0-0.1)	189 (92-369)	3 (1.5-5.7)	0 (0-1)	0 (0-0)	38 (18-73)	0.6 (0.3-1.2)
Paraguay	Gonorrhoea and chlamydia			47 (21-96)	0.6 (0.3-1.3)			5 (1-11)	0.1 (0-0.1)
Paraguay	Intra-abdominal infections	584 (361-884)	10.7 (6.7-16.2)	14,900 (8,800-23,600)	253.6 (150.7-396.2)	152 (93-237)	2.8 (1.7-4.3)	3,910 (2,230-6,410)	66.4 (38.3-107.8)
Paraguay	LRI and thorax infections	1,040 (710-1,480)	19.2 (13.2-27.4)	28,800 (18,800-42,200)	488.9 (321.1-716.5)	266 (177-387)	4.9 (3.3-7.2)	7,250 (4,660-10,700)	123.4 (79.4-182.4)
Paraguay	TB	26 (8-68)	0.4 (0.1-1.2)	953 (297-2,490)	14.8 (4.6-38.8)	12 (2-42)	0.2 (0-0.7)	427 (60-1,450)	6.7 (0.9-22.8)
Paraguay	Typhoid, paratyphoid, and iNTS	1 (0-2)	0 (0-0)	36 (10-89)	0.5 (0.2-1.3)	0 (0-0)	0 (0-0)	7 (1-22)	0.1 (0-0.3)
Paraguay	UTI	242 (152-350)	4.6 (2.9-6.7)	4,940 (3,170-7,360)	87 (55.7-128.9)	59 (36-88)	1.1 (0.7-1.7)	1,190 (742-1,820)	21 (12.9-32.2)
Peru	All	20,900 (14,600-29,200)	64 (44.7-89.6)	567,000 (377,000-822,000)	1757 (1170.8-2545)	5,220 (3,580-7,420)	16 (11-22.8)	140,000 (92,500-205,000)	434.5 (286.6-634)
Peru	Bacterial skin infections	276 (98-625)	0.8 (0.3-1.9)	6,140 (2,260-13,100)	18.8 (6.8-40.6)	64 (22-146)	0.2 (0.1-0.5)	1,410 (512-3,080)	4.3 (1.6-9.5)
Peru	Bone and joint infections	38 (10-88)	0.1 (0-0.3)	827 (215-2,050)	2.5 (0.7-6.3)	10 (3-23)	0 (0-0.1)	209 (53-541)	0.6 (0.2-1.7)
Peru	BSI	4,570 (2,440-7,430)	14.2 (7.6-23)	165,000 (99,300-255,000)	519.2 (314-794.9)	1,150 (629-1,930)	3.6 (2-6)	41,300 (24,500-64,700)	129.9 (77.6-202.4)
Peru	Cardiac infections	123 (86-172)	0.4 (0.3-0.5)	3,700 (2,510-5,410)	11.1 (7.5-16.3)	31 (22-44)	0.1 (0.1-0.1)	924 (618-1,350)	2.8 (1.9-4)
Peru	CNS infections	83 (50-144)	0.3 (0.2-0.4)	4,190 (2,480-6,980)	12.7 (7.4-21.1)	19 (11-32)	0.1 (0-0.1)	931 (541-1,570)	2.8 (1.6-4.8)
Peru	Diarrhoea	15 (7-30)	0 (0-0.1)	1,830 (573-4,880)	5.7 (1.8-15.2)	3 (1-7)	0 (0-0)	285 (108-635)	0.9 (0.3-2)
Peru	Gonorrhoea and chlamydia			147 (60-319)	0.4 (0.2-0.9)			16 (4-41)	0 (0-0.1)
Peru	Intra-abdominal infections	3,650 (2,170-5,580)	11.3 (6.7-17.3)	91,100 (52,300-144,000)	278.8 (160.5-439.3)	927 (542-1,480)	2.9 (1.7-4.6)	23,200 (13,200-37,800)	71 (40.5-115.1)

Peru	LRI and thorax infections	10,400 (7,740-13,600)	31.5 (23.5-41.6)	252,000 (177,000-348,000)	781.3 (545.1-1079.2)	2,480 (1,760-3,430)	7.5 (5.3-10.5)	58,700 (39,700-82,800)	181.8 (122.8-257)
Peru	TB	457 (235-839)	1.4 (0.7-2.5)	17,300 (8,890-30,700)	50.7 (26.1-90)	218 (43-522)	0.7 (0.1-1.6)	7,470 (1,480-17,600)	22 (4.4-51.6)
Peru	Typhoid, paratyphoid, and iNTS	7 (3-13)	0 (0-0)	393 (151-800)	1.2 (0.5-2.4)	1 (0-3)	0 (0-0)	81 (15-206)	0.2 (0-0.6)
Peru	UTI	1,300 (875-1,830)	4 (2.7-5.6)	24,400 (16,100-35,100)	74.8 (49.4-107.5)	316 (208-452)	1 (0.6-1.4)	5,900 (3,850-8,620)	18.1 (11.8-26.5)
Saint Kitts and Nevis	All	39 (27-54)	70 (48.4-97.6)	1,010 (677-1,470)	1691.2 (1137.3-2441.9)	10 (7-14)	18 (12.1-25.8)	260 (169-384)	432.3 (281.6-637.3)
Saint Kitts and Nevis	Bacterial skin infections	3 (1-4)	5 (2.9-8.5)	58 (31-106)	95.4 (53-168)	1 (0-1)	1.1 (0.6-2)	13 (7-24)	21.5 (11.3-39.1)
Saint Kitts and Nevis	Bone and joint infections	0 (0-0)	0.3 (0.1-0.6)	4 (1-8)	5.4 (1.5-12.8)	0 (0-0)	0.1 (0-0.2)	1 (0-2)	1.4 (0.4-3.4)
Saint Kitts and Nevis	BSI	13 (7-21)	23.6 (13-38.4)	367 (209-580)	616.3 (348.8-970.6)	4 (2-6)	6.4 (3.4-10.4)	99 (55-157)	165.3 (91.8-262)
Saint Kitts and Nevis	Cardiac infections	0 (0-1)	0.8 (0.5-1.1)	12 (7-18)	18.6 (11.8-27.9)	0 (0-0)	0.2 (0.1-0.3)	3 (2-5)	5 (3.1-7.5)
Saint Kitts and Nevis	CNS infections	0 (0-0)	0.5 (0.3-0.8)	12 (7-20)	22 (13.4-37.1)	0 (0-0)	0.1 (0.1-0.2)	3 (2-5)	5.1 (3-8.5)
Saint Kitts and Nevis	Diarrhoea	0 (0-0)	0 (0-0.1)	1 (1-3)	3 (1.5-5.5)	0 (0-0)	0 (0-0)	0 (0-0)	0.6 (0.3-1.1)
Saint Kitts and Nevis	Gonorrhoea and chlamydia			1 (0-2)	0.9 (0.2-2.9)			0 (0-0)	0.1 (0-0.3)
Saint Kitts and Nevis	Intra-abdominal infections	8 (5-12)	12.7 (8.1-18.6)	203 (121-314)	298.5 (179.7-456.1)	2 (1-3)	3.3 (2-4.8)	52 (30-81)	76.5 (44.5-119.1)
Saint Kitts and Nevis	LRI and thorax infections	10 (7-14)	19.6 (14.3-26.6)	264 (180-375)	485.6 (336.3-673.3)	3 (2-4)	5 (3.5-6.9)	66 (43-96)	121 (81-173.3)
Saint Kitts and Nevis	TB	0 (0-0)	0 (0-0.1)	1 (0-3)	1.1 (0.4-3.8)	0 (0-0)	0 (0-0.1)	0 (0-1)	0.4 (0.1-1.8)

Saint Kitts and Nevis	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0.1 (0-0.1)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Saint Kitts and Nevis	UTI	4 (3-5)	7.5 (5.3-10)	89 (62-124)	144.1 (100.5-197.7)	1 (1-1)	1.8 (1.3-2.5)	22 (15-32)	35.3 (24.2-49.7)
Saint Lucia	All	111 (74-159)	57 (37.8-81.8)	2,610 (1,660-3,830)	1462.9 (930.1-2161.3)	29 (19-42)	14.8 (9.6-21.6)	675 (426-1,010)	376.2 (236.8-565.6)
Saint Lucia	Bacterial skin infections	8 (5-14)	4.1 (2.4-7)	156 (89-276)	77.3 (44.6-135.5)	2 (1-3)	0.9 (0.5-1.6)	36 (19-63)	17.9 (9.7-30.9)
Saint Lucia	Bone and joint infections	0 (0-1)	0.2 (0.1-0.5)	9 (2-23)	4.6 (1.2-11.2)	0 (0-0)	0.1 (0-0.1)	2 (1-6)	1.2 (0.3-3.1)
Saint Lucia	BSI	36 (19-62)	18.9 (9.8-32.2)	892 (466-1,460)	524.9 (291.9-842.2)	10 (5-17)	5.1 (2.6-8.6)	238 (126-393)	139.2 (76.3-226.5)
Saint Lucia	Cardiac infections	4 (3-5)	1.8 (1.3-2.3)	93 (71-123)	45.3 (34.3-60.1)	1 (1-1)	0.5 (0.4-0.6)	25 (19-33)	12.1 (9.1-16.1)
Saint Lucia	CNS infections	1 (1-1)	0.5 (0.3-0.8)	36 (23-60)	23.6 (14.9-38.5)	0 (0-0)	0.1 (0.1-0.2)	8 (5-14)	5.5 (3.4-9)
Saint Lucia	Diarrhoea	0 (0-0)	0 (0-0)	2 (1-3)	1.3 (0.7-2.3)	0 (0-0)	0 (0-0)	0 (0-1)	0.3 (0.1-0.5)
Saint Lucia	Gonorrhoea and chlamydia			2 (1-3)	0.9 (0.4-1.7)			0 (0-0)	0.1 (0-0.2)
Saint Lucia	Intra-abdominal infections	19 (12-29)	9.3 (5.8-14)	467 (281-732)	223.7 (133.9-350.4)	5 (3-8)	2.4 (1.5-3.8)	122 (71-199)	58.5 (34.1-95.1)
Saint Lucia	LRI and thorax infections	31 (21-45)	16.7 (11.4-23.6)	733 (486-1,070)	454.9 (294.3-671.9)	8 (5-12)	4.3 (2.9-6.3)	188 (121-283)	114.5 (72-171.3)
Saint Lucia	TB	0 (0-0)	0 (0-0.1)	2 (1-8)	1.2 (0.4-3.9)	0 (0-0)	0 (0-0.1)	1 (0-4)	0.5 (0.1-2.2)
Saint Lucia	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0.1 (0-0.1)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Saint Lucia	UTI	11 (8-17)	5.6 (3.8-8.3)	217 (144-324)	105.3 (70.3-156.2)	3 (2-4)	1.4 (0.9-2.1)	54 (36-83)	26.3 (17.3-40.1)
Saint Vincent and the Grenadines	All	80 (54-112)	66.4 (44.8-93.5)	1,990 (1,310-2,840)	1738.4 (1145.9-2490.3)	20 (14-30)	17 (11.3-24.7)	507 (326-745)	440.9 (282.3-642.7)

Saint Vincent and the Grenadines	Bacterial skin infections	6 (3-10)	4.8 (2.7-8.5)	121 (66-210)	96 (53.1-166.5)	1 (1-2)	1.1 (0.6-1.9)	28 (14-48)	22 (11.6-38.2)
Saint Vincent and the Grenadines	Bone and joint infections	0 (0-1)	0.3 (0.1-0.7)	7 (2-17)	5.5 (1.4-13.3)	0 (0-0)	0.1 (0-0.2)	2 (0-5)	1.4 (0.4-3.6)
Saint Vincent and the Grenadines	BSI	26 (14-44)	22 (12-36.5)	702 (397-1,100)	639.9 (372.5-984.7)	7 (4-12)	5.8 (3.1-9.7)	183 (105-291)	165.9 (97.8-260.4)
Saint Vincent and the Grenadines	Cardiac infections	1 (1-1)	0.8 (0.5-1.1)	26 (18-35)	20.3 (14.4-28.1)	0 (0-0)	0.2 (0.1-0.3)	7 (5-9)	5.3 (3.7-7.4)
Saint Vincent and the Grenadines	CNS infections	1 (0-1)	0.6 (0.4-1)	31 (20-50)	30.3 (19.1-48.9)	0 (0-0)	0.1 (0.1-0.2)	7 (4-12)	7.1 (4.3-11.7)
Saint Vincent and the Grenadines	Diarrhoea	0 (0-0)	0 (0-0.1)	2 (1-3)	2.1 (1.2-3.5)	0 (0-0)	0 (0-0)	0 (0-1)	0.5 (0.3-0.8)
Saint Vincent and the Grenadines	Gonorrhoea and chlamydia			1 (1-2)	0.9 (0.4-1.7)			0 (0-0)	0.1 (0-0.2)
Saint Vincent and the Grenadines	Intra-abdominal infections	13 (8-20)	10.2 (6.4-15.5)	329 (197-510)	250.6 (149.9-390.7)	3 (2-5)	2.6 (1.6-4.1)	85 (50-136)	64.7 (37.8-104.7)
Saint Vincent and the Grenadines	LRI and thorax infections	25 (17-36)	21.2 (14.5-30.3)	617 (413-895)	570.1 (374.2-829.8)	6 (4-9)	5.5 (3.6-8)	157 (102-234)	143.7 (93-211.7)

Saint Vincent and the Grenadines	TB	0 (0-0)	0 (0-0.1)	2 (1-6)	1.4 (0.4-4.4)	0 (0-0)	0 (0-0.1)	1 (0-3)	0.6 (0.1-2.5)
Saint Vincent and the Grenadines	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0.1 (0-0.2)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Saint Vincent and the Grenadines	UTI	8 (6-10)	6.4 (4.8-8.7)	154 (114-211)	121.3 (89.7-164.9)	2 (1-3)	1.6 (1.1-2.2)	38 (27-53)	29.7 (21.4-41.5)
Suriname	All	390 (263-545)	71.6 (48.5-99.6)	11,800 (7,810-17,000)	2225.8 (1477.3-3198.5)	98 (65-141)	18.1 (12-25.6)	2,950 (1,910-4,320)	555.1 (359.7-805.2)
Suriname	Bacterial skin infections	28 (17-45)	5.1 (3.2-8.1)	638 (407-1,020)	111.5 (71.5-177.3)	6 (4-10)	1.1 (0.7-1.9)	146 (86-235)	25.5 (15.2-40.8)
Suriname	Bone and joint infections	2 (0-4)	0.3 (0.1-0.7)	37 (10-87)	6.3 (1.7-14.9)	0 (0-1)	0.1 (0-0.2)	9 (3-23)	1.6 (0.4-3.9)
Suriname	BSI	121 (67-194)	22.4 (12.6-35.6)	4,160 (2,510-6,380)	812.4 (500.5-1237.3)	31 (17-50)	5.8 (3.2-9.2)	1,060 (633-1,660)	207.4 (126.8-320)
Suriname	Cardiac infections	3 (2-4)	0.5 (0.4-0.7)	89 (62-122)	15.1 (10.6-20.6)	1 (1-1)	0.1 (0.1-0.2)	23 (16-32)	3.9 (2.7-5.4)
Suriname	CNS infections	4 (3-8)	0.8 (0.5-1.4)	239 (140-394)	46.4 (27.1-76.3)	1 (1-2)	0.2 (0.1-0.3)	56 (32-94)	10.8 (6.2-18.3)
Suriname	Diarrhoea	0 (0-0)	0.1 (0-0.1)	16 (8-29)	3.3 (1.7-6.1)	0 (0-0)	0 (0-0)	4 (2-6)	0.7 (0.4-1.4)
Suriname	Gonorrhoea and chlamydia			6 (2-12)	1 (0.4-2)			1 (0-1)	0.1 (0-0.2)
Suriname	Intra-abdominal infections	82 (53-120)	14.3 (9.2-21)	2,100 (1,280-3,170)	352.1 (216.9-529.9)	21 (13-31)	3.6 (2.3-5.3)	532 (324-821)	89.3 (54.7-137.3)
Suriname	LRI and thorax infections	111 (75-159)	21 (14.3-29.9)	3,690 (2,430-5,310)	732 (477.7-1064)	29 (19-42)	5.4 (3.5-7.9)	917 (587-1,350)	180.8 (115-264)
Suriname	TB	0 (0-0)	0 (0-0.1)	6 (2-17)	1 (0.3-2.9)	0 (0-0)	0 (0-0)	2 (0-9)	0.4 (0-1.5)
Suriname	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	2 (1-5)	0.4 (0.2-0.9)	0 (0-0)	0 (0-0)	0 (0-1)	0.1 (0-0.2)

Suriname	UTI	39 (26-53)	7 (4.8-9.5)	836 (568-1,150)	144.4 (98-197.5)	9 (6-13)	1.7 (1.1-2.4)	201 (133-282)	34.6 (23.1-48.5)
The Bahamas	All	217 (149-309)	62.6 (43.7-88.3)	5,830 (3,920-8,470)	1637.9 (1121.7-2349.3)	52 (34-75)	14.9 (10-21.4)	1,390 (910-2,070)	388.1 (256.9-575.6)
The Bahamas	Bacterial skin infections	13 (8-22)	3.8 (2.3-6.5)	280 (164-475)	75 (44.3-125.9)	3 (1-5)	0.8 (0.4-1.5)	60 (33-103)	16.1 (8.9-28)
The Bahamas	Bone and joint infections	1 (0-2)	0.2 (0.1-0.6)	21 (6-51)	5.5 (1.5-13)	0 (0-0)	0.1 (0-0.1)	5 (1-12)	1.2 (0.3-3)
The Bahamas	BSI	64 (35-106)	18.5 (10.1-30.5)	1,880 (1,070-3,100)	553.4 (324.1-878.5)	16 (8-26)	4.5 (2.4-7.4)	452 (256-751)	133 (77.8-214.8)
The Bahamas	Cardiac infections	4 (3-6)	1.1 (0.9-1.5)	139 (101-186)	34.1 (25-45.5)	1 (1-1)	0.3 (0.2-0.4)	33 (24-46)	8.2 (5.9-11.2)
The Bahamas	CNS infections	2 (1-3)	0.5 (0.3-0.9)	82 (50-142)	24.4 (14.7-41.8)	0 (0-1)	0.1 (0.1-0.2)	18 (11-32)	5.3 (3.2-9.4)
The Bahamas	Diarrhoea	0 (0-0)	0 (0-0)	4 (2-7)	1.4 (0.7-2.5)	0 (0-0)	0 (0-0)	1 (0-1)	0.3 (0.1-0.4)
The Bahamas	Gonorrhoea and chlamydia			3 (1-6)	0.8 (0.3-1.6)			0 (0-1)	0.1 (0-0.2)
The Bahamas	Intra-abdominal infections	39 (24-59)	10.4 (6.5-15.7)	1,060 (636-1,660)	263.7 (159.4-410.8)	9 (6-15)	2.5 (1.5-3.9)	259 (151-416)	64.1 (37.8-102.3)
The Bahamas	LRI and thorax infections	72 (52-101)	21.7 (15.9-29.9)	1,860 (1,310-2,640)	548 (395.9-761.1)	17 (12-25)	5.1 (3.6-7.3)	435 (293-633)	127.7 (86.8-182.9)
The Bahamas	TB	1 (0-2)	0.2 (0.1-0.4)	34 (15-66)	8.3 (3.7-16.1)	0 (0-1)	0.1 (0-0.2)	12 (3-32)	2.9 (0.8-7.6)
The Bahamas	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0.1)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
The Bahamas	UTI	21 (15-29)	6.1 (4.5-8.4)	468 (335-666)	123.3 (89.1-175.5)	5 (4-7)	1.4 (1-2)	111 (78-163)	29.3 (20.8-42.5)
Trinidad and Tobago	All	911 (558-1,400)	54.4 (33.3-83.6)	22,200 (13,200-35,100)	1492.5 (887.4-2360.9)	232 (138-363)	13.8 (8.3-21.6)	5,620 (3,260-8,940)	376.3 (219.8-592.7)
Trinidad and Tobago	Bacterial skin infections	51 (23-99)	2.9 (1.3-5.6)	1,040 (451-2,020)	59.3 (26.5-114.5)	12 (5-24)	0.7 (0.3-1.4)	242 (99-497)	13.8 (5.8-28)
Trinidad and Tobago	Bone and joint infections	4 (1-10)	0.2 (0.1-0.5)	85 (21-217)	5 (1.3-12.5)	1 (0-3)	0.1 (0-0.1)	22 (6-60)	1.3 (0.3-3.4)

Trinidad and Tobago	BSI	328 (160-563)	19.8 (9.8-33.6)	8,430 (4,350-14,300)	594.9 (316.3-978.7)	84 (41-146)	5 (2.5-8.7)	2,140 (1,090-3,670)	150.6 (79.9-248.5)
Trinidad and Tobago	Cardiac infections	14 (8-20)	0.8 (0.5-1.2)	347 (211-522)	21 (12.9-31.7)	4 (2-5)	0.2 (0.1-0.3)	88 (52-134)	5.3 (3.1-8.1)
Trinidad and Tobago	CNS infections	6 (4-10)	0.4 (0.2-0.6)	232 (143-392)	17.5 (10.7-29.2)	1 (1-2)	0.1 (0.1-0.2)	55 (32-92)	4.1 (2.4-7)
Trinidad and Tobago	Diarrhoea	0 (0-0)	0 (0-0)	16 (8-29)	1.3 (0.7-2.4)	0 (0-0)	0 (0-0)	3 (1-5)	0.3 (0.1-0.5)
Trinidad and Tobago	Gonorrhoea and chlamydia			10 (4-20)	0.8 (0.3-1.6)			1 (0-2)	0.1 (0-0.2)
Trinidad and Tobago	Intra-abdominal infections	157 (92-255)	8.8 (5.2-14.4)	3,770 (2,140-6,240)	217.8 (121.8-363.9)	40 (23-66)	2.3 (1.3-3.7)	968 (536-1,630)	56 (31-94.7)
Trinidad and Tobago	LRI and thorax infections	270 (170-409)	16.7 (10.6-25.2)	6,530 (4,030-9,960)	479 (295.3-729.1)	70 (42-109)	4.3 (2.6-6.7)	1,670 (989-2,620)	121.1 (72.6-186.8)
Trinidad and Tobago	TB	0 (0-1)	0 (0-0.1)	14 (5-42)	0.8 (0.3-2.5)	0 (0-1)	0 (0-0)	5 (1-21)	0.3 (0-1.2)
Trinidad and Tobago	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Trinidad and Tobago	UTI	82 (49-127)	4.7 (2.8-7.2)	1,680 (992-2,640)	95.1 (56.4-150.1)	20 (12-32)	1.1 (0.7-1.8)	414 (240-657)	23.4 (13.6-36.8)
United States	All	173,000 (120,000-243,000)	30.7 (21-43.4)	3,390,000 (2,260,000-4,860,000)	721.4 (474.3-1045.9)	41,900 (28,300-59,600)	7.4 (5-10.7)	821,000 (535,000-1,190,000)	174.1 (111.7-257)
United States	Bacterial skin infections	5,730 (2,700-11,300)	1 (0.5-2)	108,000 (51,600-212,000)	22 (10.5-43.1)	1,100 (467-2,290)	0.2 (0.1-0.4)	20,700 (8,970-43,000)	4.2 (1.9-8.7)
United States	Bone and joint infections	762 (250-1,750)	0.1 (0-0.3)	15,600 (4,850-36,000)	3.2 (0.9-7.4)	162 (49-377)	0 (0-0.1)	3,280 (977-7,900)	0.7 (0.2-1.6)
United States	BSI	60,800 (32,500-102,000)	11.2 (6.1-18.7)	1,300,000 (712,000-2,140,000)	290.6 (158.8-470.5)	15,000 (7,710-25,200)	2.8 (1.4-4.6)	319,000 (171,000-522,000)	70.7 (38.2-116)
United States	Cardiac infections	6,650 (4,130-8,890)	1.2 (0.7-1.5)	123,000 (79,600-166,000)	25.5 (16.6-34.9)	1,640 (1,000-2,260)	0.3 (0.2-0.4)	29,800 (18,800-41,200)	6.2 (3.9-8.6)

United States	CNS infections	536 (344-953)	0.1 (0.1-0.2)	18,700 (12,200-32,000)	5.7 (3.8-9.2)	111 (67-205)	0 (0-0)	3,830 (2,360-6,780)	1.2 (0.7-1.9)
United States	Diarrhoea	142 (77-237)	0 (0-0)	4,740 (2,480-8,150)	1.3 (0.7-2.3)	31 (15-54)	0 (0-0)	856 (418-1,500)	0.2 (0.1-0.4)
United States	Gonorrhoea and chlamydia			230 (134-361)	0.1 (0-0.1)			24 (7-49)	0 (0-0)
United States	Intra-abdominal infections	35,600 (23,000-51,700)	6.5 (4.2-9.6)	781,000 (499,000-1,150,000)	163.5 (102.7-243.2)	9,080 (5,820-13,100)	1.7 (1.1-2.4)	199,000 (126,000-290,000)	41.7 (25.7-61.5)
United States	LRI and thorax infections	46,500 (35,100-62,300)	7.9 (5.9-10.6)	788,000 (573,000-1,100,000)	163.5 (119-227.9)	11,000 (7,610-15,500)	1.9 (1.3-2.7)	186,000 (125,000-269,000)	38.4 (26.2-55.8)
United States	TB	129 (91-184)	0 (0-0)	3,190 (2,340-4,490)	0.7 (0.5-1)	36 (1-86)	0 (0-0)	782 (33-1,870)	0.2 (0-0.4)
United States	Typhoid, paratyphoid, and iNTS	2 (1-2)	0 (0-0)	53 (35-65)	0 (0-0)	0 (0-1)	0 (0-0)	10 (2-18)	0 (0-0)
United States	UTI	16,000 (12,600-21,000)	2.6 (2-3.4)	243,000 (185,000-334,000)	45.5 (34-63.3)	3,790 (2,890-5,090)	0.6 (0.5-0.8)	57,900 (42,300-81,900)	10.8 (7.8-15.5)
Uruguay	All Bacterial	2,790 (2,050-3,750)	47.4 (34.4-63.9)	48,100 (34,100-65,800)	1023.2 (714-1415.9)	692 (496-952)	11.8 (8.4-16.3)	11,900 (8,320-16,800)	254.2 (172.7-361.9)
Uruguay	skin infections	92 (52-161)	1.5 (0.8-2.6)	1,360 (743-2,460)	25.4 (13.9-47.2)	21 (11-39)	0.3 (0.2-0.6)	305 (155-571)	5.7 (2.9-10.8)
Uruguay	Bone and joint infections	6 (2-15)	0.1 (0-0.3)	108 (31-260)	2.2 (0.6-5.2)	1 (0-3)	0 (0-0.1)	25 (7-59)	0.5 (0.1-1.2)
Uruguay	BSI	735 (389-1,240)	13.4 (7.3-22.6)	15,100 (8,290-24,600)	353.4 (199.4-557.2)	187 (98-318)	3.4 (1.8-5.7)	3,810 (2,060-6,330)	89.5 (49.2-144)
Uruguay	Cardiac infections	115 (76-149)	1.9 (1.3-2.5)	1,940 (1,350-2,470)	37.8 (26.8-48.1)	29 (19-39)	0.5 (0.3-0.6)	491 (335-642)	9.5 (6.6-12.3)
Uruguay	CNS infections	8 (5-14)	0.2 (0.1-0.3)	231 (137-438)	6.3 (3.6-12)	2 (1-3)	0 (0-0.1)	54 (30-104)	1.5 (0.8-2.8)
Uruguay	Diarrhoea	1 (1-2)	0 (0-0)	55 (31-88)	1.7 (0.9-2.7)	0 (0-1)	0 (0-0)	11 (6-18)	0.3 (0.2-0.6)
Uruguay	Gonorrhoea and chlamydia			16 (8-28)	0.5 (0.3-0.8)			2 (0-3)	0 (0-0.1)
Uruguay	Intra-abdominal infections	548 (375-767)	9.4 (6.4-13.2)	9,560 (6,420-13,700)	189.4 (124.6-273.9)	137 (94-194)	2.3 (1.6-3.3)	2,390 (1,600-3,440)	47.2 (31.2-69.2)
Uruguay	LRI and thorax infections	869 (676-1,130)	14.4 (11.1-19.1)	14,100 (10,500-19,100)	305.3 (223.7-413.5)	215 (157-292)	3.6 (2.6-4.9)	3,500 (2,480-4,870)	75.4 (52.8-106.4)

Uruguay	TB	1 (1-2)	0 (0-0)	33 (19-63)	0.8 (0.4-1.5)	0 (0-1)	0 (0-0)	11 (2-31)	0.2 (0-0.7)
Uruguay	Typhoid, paratyphoid, and iNTS	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Uruguay	UTI	417 (290-526)	6.4 (4.5-8.2)	5,610 (3,980-7,290)	100.4 (73.1-132.2)	100 (70-130)	1.6 (1.1-2)	1,350 (955-1,810)	24.2 (17-33.1)
Venezuela	All	12,000 (7,460-18,300)	43.8 (27.5-66.4)	358,000 (221,000-550,000)	1324.2 (822.5-2018.8)	3,050 (1,870-4,700)	11.1 (6.9-17.1)	89,900 (55,100-138,000)	331.5 (204-507)
Venezuela	Bacterial skin infections	294 (70-728)	1.1 (0.3-2.6)	6,750 (1,840-16,000)	23.5 (6.4-56.2)	69 (16-174)	0.2 (0.1-0.6)	1,570 (407-3,780)	5.4 (1.4-13.2)
Venezuela	Bone and joint infections	41 (11-100)	0.1 (0-0.4)	1,060 (282-2,680)	3.6 (1-9.1)	10 (3-26)	0 (0-0.1)	272 (70-709)	0.9 (0.2-2.4)
Venezuela	BSI	4,470 (2,240-7,530)	16.4 (8.3-27.5)	140,000 (76,300-226,000)	524.4 (290.6-839.2)	1,180 (604-2,020)	4.3 (2.2-7.4)	36,600 (20,100-60,500)	136.9 (77-222.6)
Venezuela	Cardiac infections	123 (73-191)	0.4 (0.3-0.7)	3,310 (1,990-5,050)	11.4 (6.9-17.4)	32 (19-51)	0.1 (0.1-0.2)	872 (521-1,360)	3 (1.8-4.7)
Venezuela	CNS infections	111 (68-185)	0.4 (0.2-0.7)	5,870 (3,650-9,450)	22.2 (13.8-35.4)	26 (16-44)	0.1 (0.1-0.2)	1,360 (829-2,200)	5.1 (3.2-8.2)
Venezuela	Diarrhoea	39 (20-70)	0.2 (0.1-0.3)	2,460 (1,260-4,370)	10 (5.1-17.9)	9 (3-18)	0 (0-0.1)	482 (177-1,030)	2 (0.7-4.2)
Venezuela	Gonorrhoea and chlamydia			115 (50-240)	0.4 (0.2-0.9)			11 (2-26)	0 (0-0.1)
Venezuela	Intra-abdominal infections	2,530 (1,520-3,920)	8.9 (5.4-13.8)	65,000 (37,700-103,000)	221.5 (129.6-353.4)	640 (380-1,040)	2.3 (1.4-3.7)	16,500 (9,400-27,300)	56.2 (32.1-93.2)
Venezuela	LRI and thorax infections	3,440 (2,300-5,050)	12.8 (8.6-18.8)	112,000 (74,400-166,000)	431.8 (285-634.9)	841 (534-1,260)	3.1 (2-4.7)	26,600 (16,800-40,000)	102.2 (64.9-153.1)
Venezuela	TB	74 (32-168)	0.3 (0.1-0.6)	2,430 (1,060-5,460)	8.3 (3.7-18.8)	30 (5-97)	0.1 (0-0.3)	885 (161-2,870)	3 (0.6-9.8)
Venezuela	Typhoid, paratyphoid, and iNTS	1 (0-2)	0 (0-0)	14 (1-42)	0 (0-0.1)	0 (0-0)	0 (0-0)	3 (0-9)	0 (0-0)
Venezuela	UTI	871 (507-1,410)	3.2 (1.9-5.1)	19,100 (10,800-31,200)	66.9 (38.4-109.1)	216 (123-354)	0.8 (0.4-1.3)	4,740 (2,640-7,890)	16.6 (9.3-27.5)

Abbreviations: BSI = Bloodstream Infection, CNS = Central Nervous System, DALYs = Disability Adjusted Life Years, iNTS = invasive Non-typhoidal Salmonella, LRI = Lower Respiratory Infection, TB = tuberculosis, UTI = Urinary Tract Infection, WHO = World Health Organization