

Supplementary information

Causes and consequences of child growth faltering in low-resource settings

In the format provided by the authors and unedited

Supplement Information for:

Causes and consequences of child growth faltering in low- and middle-income countries

Andrew Mertens,^{1*} Jade Benjamin-Chung,^{2,3,4} John M Colford Jr,¹ Jeremy Coyle¹, Mark J van der Laan,¹ Alan E Hubbard,¹ Sonali Rosete,¹ Ivana Malenica,¹ Nima Hejazi,¹ Oleg Sofrygin,¹ Wilson Cai,¹ Haodong Li,¹ Anna Nguyen,¹ Nolan N Pokpongkiat,¹ Stephanie Djajadi,¹ Anmol Seth,¹ Esther Jung,¹ Esther O Chung,¹ Wendy Jilek,¹ Vishak Subramoney,⁵ Ryan Hafen,⁶ Jonas Häggström,⁷ Thea Norman,⁸ Kenneth H Brown,⁹ Parul Christian,¹⁰ Benjamin F. Arnold,¹¹ and members of the *ki* Child Growth Consortium

¹ Division of Epidemiology & Biostatistics, University of California, Berkeley, 2121 Berkeley Way Rm 5302 Berkeley, CA 94720-7360

² Department of Epidemiology & Population Health, Stanford University, Stanford University, 300 Pasteur Dr., Stanford, CA 94305

³ Division of Epidemiology & Biostatistics, University of California, Berkeley, 2121 Berkeley Way Rm 5302 Berkeley, CA 94720-7360

⁴ Chan Zuckerberg Biohub, San Francisco, CA 94158

⁵ DVPL Tech

⁶ Hafen Consulting, LLC, West Richland WA, 99353

⁷ Cytel Inc, 1050 Winter St Suite 2700 Waltham, MA 02451, USA

⁸ Quantitative Sciences, Bill & Melinda Gates Foundation, 500 5th Ave N, Seattle, WA 98109

⁹ Department of Nutrition, University of California, Davis, 3135 Meyer Hall, Davis, CA 95616-5270

¹⁰ Center for Human Nutrition, Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD 21205

¹¹ Francis I. Proctor Foundation and Department of Ophthalmology, University of California, San Francisco, 490 Illinois St, San Francisco, CA 94158

Corresponding authors:

Andrew Mertens, PhD, amertens@berkeley.edu, Research Data Analyst, Division of Epidemiology & Biostatistics, University of California, Berkeley, 2121 Berkeley Way Rm 5302 Berkeley, CA 94720-7360

Benjamin F. Arnold, PhD, ben.arnold@ucsf.edu, Associate Professor, Francis I. Proctor Foundation, University of California, San Francisco, 490 Illinois St, San Francisco, CA 94158

Table of Contents

Supplementary Note 1. Directed Acyclic Diagrams (DAGs)	2
1.1 Overview	2
1.2 Simplified example DAG:.....	3
1.3 DAG used: Association between maternal weight and child growth faltering	4
1.4 Table of Adjustment Covariates	4
Supplementary Note 2. Complete-Case Sensitivity Analyses	9
2.1 Heatmap of covariate missingness by study	9
2.2 Sensitivity analysis to complete case versus median/mode imputed analyses.....	9
Plots comparing population intervention effect estimates (the estimates presented in Figures 2 and 3) compared to estimates only using complete-case data (excluding any observations missing covariate information).	9
Supplementary Note 3. Sensitivity analysis using fixed effects models.....	11
3.1 Overview	11
3.2 Primary manuscript figures recreated with estimates pooled using fixed effects.....	12
Supplementary Note 4. Comparison of Adjusted and Unadjusted Population Intervention Effects	17
Supplementary Note 5. Spline plots: all exposures	17
Supplementary Note 6. Sensitivity to dropping PROBIT trial.....	77
6.1 Overview	77
6.2 Comparison of attributable differences estimated with and without the PROBIT trial.....	78
Supplementary Note 7 Average treatment effect and relative risk plots	78

Supplementary Note 1. Directed Acyclic Diagrams (DAGs)


1.1 Overview

For each exposure, we used the directed acyclic graph (DAG) framework to identify potential confounders from the broader set of exposures used in the analysis. We did not adjust for characteristics that were assumed to be intermediate on the causal path between any exposure and the outcome, because while controlling for mediators may help adjust for unmeasured confounders in some conditions, it can also lead to collider bias. As DAG's were hand-drawn and are complex for the large number of exposures, we show an example DAG here and then list all potential considered for each exposure in a table below.


The DAG below shows the set of adjusted-for exposures and unadjusted-for exposures for the estimated association between maternal weight and child growth faltering. Grey variables are on the causal pathway and are not adjusted for, while blue variables were considered potential confounders and adjusted for. Note collinear variables like maternal BMI are shown as on the causal pathway in this DAG, which may not be biologically accurate, but they were not adjusted for so are included with the potential mediators.

Given the exposures measured in each study, we attempted to maximally adjust for confounding without adjusting for mediators on the casual pathway, which would remove part of the effect of the exposure on the outcome. Confounders were not measured in every cohort, so there could be residual confounding in cohort-specific estimates, and actual DAGs will look different across cohort-specific estimates for each study.

1.2 Simplified example DAG:



1.3 DAG used: Association between maternal weight and child growth faltering



1.4 Table of Adjustment Covariates

Exposure	Potential adjustment covariates
# of children <5 in HH	Treatment arm , Mother’s age , Father’s age , Mother’s education , Father’s education , HH wealth , HH food security , Mother’s height , Mother’s weight , BMI , Father’s height , Single parent , # of rooms in HH , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
# of people in HH	Treatment arm , Mother’s age , Father’s age , Mother’s education , Father’s education , HH wealth , HH food security , Mother’s height , Mother’s weight , BMI , Father’s height , Single parent , # of rooms in HH , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source

Exposure	Potential adjustment covariates
# of rooms in HH	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of people in HH , # of children <5 in HH , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Any wasting <6 mo.	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of measurement , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Birth length (cm)	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Birth order	Treatment arm , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Single parent , # of rooms in HH , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Birthweight (kg)	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Breastfed hour after birth	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Gestational age at birth, Birthweight (kg) , Birth length (cm) , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Child delivered at home	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats

Exposure	Potential adjustment covariates
	drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Clean cooking fuel usage	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH, Birth order , Treats drinking water , Improved floor , Improved sanitation , Safe water source
Diarrhea <24 mo. (% days)	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Gestational age at birth, Birthweight (kg) , Birth length (cm) , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of measurement , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Diarrhea <6 mo. (% days)	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Gestational age at birth, Birthweight (kg) , Birth length (cm) , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of measurement , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Excl/Pred breastfed <6mo.	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Gestational age at birth, Birthweight (kg) , Birth length (cm) , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of measurement , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Father's age	Treatment arm , Mother's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Father's education	Treatment arm , Mother's age , Father's age , Mother's education , HH wealth , Mother's height , Mother's weight , BMI , Father's height , Gestational age at birth, Birthweight (kg) , Birth length (cm) , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Father's height	Treatment arm , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight

Exposure	Potential adjustment covariates
	, BMI , Single parent , # of rooms in HH , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Gestational age at birth	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
HH food security	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , Vaginal birth , Child delivered at home , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
HH wealth	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , Gestational age at birth, Birthweight (kg) , Birth length (cm) , Single parent , # of people in HH , # of children <5 in HH , Birth order
Improved floor	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Birth order , Treats drinking water , Clean cooking fuel usage, Improved sanitation , Safe water source
Improved sanitation	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Safe water source
Mother's age	Treatment arm , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Mother's BMI	Treatment arm , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Mother's education	Treatment arm , Mother's age , Father's age , Father's education , HH wealth , Mother's height , Mother's weight , BMI , Father's height , Gestational age at birth, Birthweight (kg) , Birth length (cm) , Single parent , # of rooms in HH

Exposure	Potential adjustment covariates
	, # of people in HH , # of children <5 in HH , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Mother's height	Treatment arm , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Father's height , Single parent , # of rooms in HH , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Mother's weight	Treatment arm , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Persistent wasting <6 mo.	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Vaginal birth , Child delivered at home , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of measurement , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Safe water source	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation
Single parent	Treatment arm , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , # of rooms in HH , # of people in HH , # of children <5 in HH , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Small for gestational age	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Treats drinking water , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source
Treats drinking water	Treatment arm , Child sex , Mother's age , Father's age , Mother's education , Father's education , HH wealth , HH food security , Mother's height , Mother's weight , BMI , Father's height , Single parent , # of rooms in HH , # of people in HH , # of children <5 in HH , Month of Birth , Birth order , Clean cooking fuel usage, Improved floor , Improved sanitation , Safe water source




The median absolute difference in pooled Z-scores was 0.009 and the mean absolute difference was 0.022.

Supplementary Note 3. Sensitivity analysis using fixed effects models

3.1 Overview


In the primary manuscript, we compared estimates pooled across studies using random effects models, which are more conservative in the presence of heterogeneity across studies. Here, we present estimates pooled using fixed effects. Inferences about estimates from fixed effects models are restricted to only the included studies, while random effects estimates assume included studies are sampled from a distribution of hypothetical studies. Fixed effect estimates had narrow confidence intervals, and therefore more risk factors had statistically significant PIEs. The fixed effect estimates are weighted towards results from larger studies, so are more heavily influenced by the large Jivita-3, Probit, and ZVITAMBO trials. But in general, the key exposures identified from the random-effects estimates also had the strongest associations when pooled using fixed-effect estimates.

3.2 Primary manuscript figures recreated with estimates pooled using fixed effects




Heatmap of significance and direction across exposure-outcome combinations of associations estimated using fixed effects.

Attributable difference - LAZ, stratified by age




Age-stratified population attributable differences in length-for-age Z-scores estimated using fixed effects.

Attributable difference - WLZ, stratified by age




Age-stratified population attributable differences in weight-for-length Z-scores estimated using fixed effects.

Population intervention effect - LAZ, stratified by region



Region-stratified population attributable differences in length-for-age Z-scores estimated using fixed effects.


Population intervention effect - WLZ, stratified by region



Region-stratified population attributable differences in weight-for-length Z-scores estimated using fixed effects.

Supplementary Note 4. Comparison of Adjusted and Unadjusted Population Intervention Effects


Plots comparing covariate-adjusted population intervention effect estimates (the estimates presented in Figures 2 and 3) compared to estimates not adjusted for potential confounders laid out in Table 1.4.




Supplementary Note 5. Spline plots: all exposures

Spline plots of HAZ and WHZ over the first two years of children's lives, stratified by levels of risk factors. Plots were constructed using the same methods as for Figures 4a and 4b for all risk factors (details in the Methods section).


Spline curves of LAZ stratified by levels of exposure: Birth length (cm)




Spline curves of LAZ stratified by levels of exposure: Birthweight (kg)




Spline curves of LAZ stratified by levels of exposure: Clean cooking fuel usage




Spline curves of LAZ stratified by levels of exposure: Breastfed hour after birth




Spline curves of LAZ stratified by levels of exposure: Enrolled stunted




Spline curves of LAZ stratified by levels of exposure: Enrolled wasted




Spline curves of LAZ stratified by levels of exposure: Father's age




Spline curves of LAZ stratified by levels of exposure: Father's education




Spline curves of LAZ stratified by levels of exposure: Father's height




Spline curves of LAZ stratified by levels of exposure: Gestational age at birth




Spline curves of LAZ stratified by levels of exposure: Child delivered at home




Spline curves of LAZ stratified by levels of exposure: HH food security




Spline curves of LAZ stratified by levels of exposure: HH wealth




Spline curves of LAZ stratified by levels of exposure: Improved floor




Spline curves of LAZ stratified by levels of exposure: Improved sanitation




Spline curves of LAZ stratified by levels of exposure: Mother's age




Spline curves of LAZ stratified by levels of exposure: Mother's education




Spline curves of LAZ stratified by levels of exposure: # of children <5 in HH




Spline curves of LAZ stratified by levels of exposure: # of people in HH




Spline curves of LAZ stratified by levels of exposure: # of rooms in HH




Spline curves of LAZ stratified by levels of exposure: Birth order




Spline curves of LAZ stratified by levels of exposure: Diarrhea <6 mo. (% days)




Spline curves of LAZ stratified by levels of exposure: Diarrhea <24 mo. (% days)




Spline curves of LAZ stratified by levels of exposure: Persistent wasting <6 mo.




Spline curves of LAZ stratified by levels of exposure: Excl/Pred breastfed <6mo.




Spline curves of LAZ stratified by levels of exposure: Safe water source




Spline curves of LAZ stratified by levels of exposure: Sex




Spline curves of LAZ stratified by levels of exposure: Single parent




Spline curves of LAZ stratified by levels of exposure: Treats drinking water




Spline curves of LAZ stratified by levels of exposure: Vaginal birth




Spline curves of WLZ stratified by levels of exposure: Birth length (cm)




Spline curves of WLZ stratified by levels of exposure: Birthweight (kg)




Spline curves of WLZ stratified by levels of exposure: Clean cooking fuel usage




Spline curves of WLZ stratified by levels of exposure: Breastfed hour after birth




Spline curves of WLZ stratified by levels of exposure: Enrolled stunted




Spline curves of WLZ stratified by levels of exposure: Enrolled wasted




Spline curves of WLZ stratified by levels of exposure: Father's age




Spline curves of WLZ stratified by levels of exposure: Father's education




Spline curves of WLZ stratified by levels of exposure: Father's height




Spline curves of WLZ stratified by levels of exposure: Gestational age at birth




Spline curves of WLZ stratified by levels of exposure: Child delivered at home




Spline curves of WLZ stratified by levels of exposure: HH food security




Spline curves of WLZ stratified by levels of exposure: HH wealth




Spline curves of WLZ stratified by levels of exposure: Improved floor




Spline curves of WLZ stratified by levels of exposure: Improved sanitation




Spline curves of WLZ stratified by levels of exposure: Mother's age




Spline curves of WLZ stratified by levels of exposure: Mother's education




Spline curves of WLZ stratified by levels of exposure: # of children <5 in HH




Spline curves of WLZ stratified by levels of exposure: # of people in HH




Spline curves of WLZ stratified by levels of exposure: # of rooms in HH




Spline curves of WLZ stratified by levels of exposure: Birth order




Spline curves of WLZ stratified by levels of exposure: Diarrhea <6 mo. (% days)




Spline curves of WLZ stratified by levels of exposure: Diarrhea <24 mo. (% days)




Spline curves of WLZ stratified by levels of exposure: Persistent wasting <6 mo.




Spline curves of WLZ stratified by levels of exposure: Excl/Pred breastfed <6mo.




Spline curves of WLZ stratified by levels of exposure: Safe water source




Spline curves of WLZ stratified by levels of exposure: Sex




Spline curves of WLZ stratified by levels of exposure: Single parent



Spline curves of WLZ stratified by levels of exposure: Treats drinking water



Spline curves of WLZ stratified by levels of exposure: Vaginal birth




Supplementary Note 6. Sensitivity to dropping PROBIT trial

6.1 Overview

The PROBIT trial, conducted in Belarus, was the only study included conducted in Europe, as opposed to Latin America, South Asia, or sub-Saharan Africa. It was also one of the largest studies, so may have an influential effect on the pooled estimates. Because of differences in income, environmental conditions, and population anthropometry, we reconducted the primary analyses without the PROBIT trial and present the results here. We found that the exclusion of the PROBIT trial did not affect the main inferences presented in the manuscript.

6.2 Comparison of attributable differences estimated with and without the PROBIT trial

Because PROBIT was the only European study, we also conducted a sensitivity analysis as to the effect of removing PROBIT on attributable differences at 24 months (within the exposures measures during the PROBIT trial). Other than for the estimated associations between father’s height and child Z-scores at 24 months (which was measured in few other studies), PROBIT is not highly influential.



Supplementary Note 7 Average treatment effect and relative risk plots



The primary parameter presented in the “Causes and Consequences of Child Growth Failure in Low- and Middle-income Countries” publications is the population intervention effect (PIE), the change in mean population Z-score if a exposure is shifted to the lowest-risk level, which combines the increase of risk associated with an exposure with the prevalence of the exposure in the population. This page of the supporting information presents plots for pooled relative risks (RR) and average treatment effects (ATE) for each level of each exposure (all the estimates shown in “Extended Data Figure 2. Heatmap of significance and direction across exposure-outcome combinations”, but presented here with point estimates and confidence intervals). Relative risk are reported for binary CGF outcomes (stunting, wasting, underweight, severe stunting, severe wasting, and severe underweight) and ATEs (mean differences) are reported for continuous Z-score outcomes.

The primary purpose of this page is to present the pooled point estimates and confidence intervals for all of the results summarized in the Extended Data Figure 2 heatmap, and to allow the reader to compare the levels of risk across categorical exposures. The population intervention effects and attributable


fraction presented in the primary manuscript show the result when shifting the exposure distribution to

the low-risk reference levels, but the results presented here allow for comparing other exposure levels, like food secure versus moderately food insecure.


Plots are grouped and titled by exposure and growth outcome, and each panel of grouped plots visualizes the relative risk between levels of the exposure and different age-specific growth failure outcomes. Reference levels are printed in the titles after "ref:".




Outcome: LAZ
Exposure:
Clean cooking fuel usage, ref: Yes




Outcome: LAZ
Exposure:
Breastfed hour after birth, ref: Yes







Outcome: LAZ
Exposure:
Father's age, ref: >=35




Outcome: LAZ
Exposure:
Father's education, ref: High











Outcome: LAZ
Exposure:
HH wealth, ref: Q4







Outcome: LAZ
Exposure:
Improved floor, ref: Yes


















Outcome: LAZ
Exposure:
Diarrhea <6 mo. (% days), ref: [0%, 2%]




Outcome: LAZ
Exposure:
Excl/Pred breastfed <6mo., ref: Yes




Outcome: LAZ
Exposure:
Rain quartile, ref: Opposite max rain




Outcome: LAZ
Exposure:
Safe water source, ref: Yes




Outcome: LAZ
Exposure:
Sex, ref: Female





Outcome: LAZ
Exposure:
Small-for-gestational age, ref: Not SGA




Outcome: LAZ
Exposure:
Single parent, ref: 0







Outcome: WHZ
Exposure:
Clean cooking fuel usage, ref: Yes




Outcome: WHZ
Exposure:
Breastfed hour after birth, ref: Yes







Outcome: WHZ
Exposure:
Father's age, ref: ≥ 35




Outcome: WHZ
Exposure:
Father's education, ref: High









Outcome: WHZ
Exposure:
HH wealth, ref: Q4




Outcome: WHZ
Exposure:
Improved floor, ref: Yes







Outcome: WHZ
Exposure:
Improved sanitation, ref: Yes




Outcome: WHZ
Exposure:
Mother's age, ref: [20-30)











Outcome: WHZ
Exposure:
of children <5 in HH, ref: 1




Outcome: WHZ
Exposure:
of people in HH, ref: 3 or less









Outcome: WHZ
Exposure:
Sex, ref: Female




Outcome: WHZ
Exposure:
Small-for-gestational age, ref: Not SGA





Outcome: WHZ
Exposure:
Single parent, ref: 0




Outcome: Stunted
Exposure:
Birth length (cm), ref: ≥ 50 cm










Outcome: Stunted
Exposure:
Breastfed hour after birth, ref: Yes




Outcome: Stunted
Exposure:
Father's age, ref: ≥ 35









Outcome: Stunted
Exposure:
HH food security, ref: Food Secure




Outcome: Stunted
Exposure:
HH wealth, ref: Q4




Outcome: Stunted
Exposure:
Improved floor, ref: Yes




Outcome: Stunted
Exposure:
Improved sanitation, ref: Yes





Outcome: Stunted
Exposure:
Mother's age, ref: [20-30)




Outcome: Stunted
Exposure:
Mother's BMI, ref: >=20 kg/m²








Outcome: Stunted
Exposure:
Mother's weight, ref: ≥ 45 kg




Outcome: Stunted
Exposure:
of children <5 in HH, ref: 1







Outcome: Stunted
Exposure:
Birth order, ref: 1




Outcome: Stunted
Exposure:
Diarrhea <6 mo. (% days), ref: [0%, 2%]





Outcome: Stunted
Exposure:
Persistent wasting <6 mo., ref: No




Outcome: Stunted
Exposure:
Excl/Pred breastfed <6mo., ref: Yes







Outcome: Stunted
Exposure:
Sex, ref: Female




Outcome: Stunted
Exposure:
Small-for-gestational age, ref: Not SGA





Outcome: Stunted
Exposure:
Single parent, ref: 0




Outcome: Ever Stunted
Exposure:
Birthweight (kg), ref: ≥ 2500 g







Outcome: Ever Stunted
Exposure:
Breastfed hour after birth, ref: Yes




Outcome: Ever Stunted
Exposure:
Father's age, ref: >=35





Outcome: Ever Stunted
Exposure:
Father's education, ref: High




Outcome: Ever Stunted
Exposure:
Father's height, ref: ≥ 162 cm







Outcome: Ever Stunted
Exposure:
HH food security, ref: Food Secure




Outcome: Ever Stunted
Exposure:
HH wealth, ref: Q4




Outcome: Ever Stunted
Exposure:
Improved floor, ref: Yes




Outcome: Ever Stunted
Exposure:
Improved sanitation, ref: Yes





Outcome: Ever Stunted
Exposure:
Mother's age, ref: [20-30)




Outcome: Ever Stunted
Exposure:
Mother's BMI, ref: ≥ 20 kg/m²








Outcome: Ever Stunted
Exposure:
Mother's weight, ref: ≥ 45 kg




Outcome: Ever Stunted
Exposure:
of children <5 in HH, ref: 1







Outcome: Ever Stunted
Exposure:
Birth order, ref: 1




Outcome: Ever Stunted
Exposure:
Safe water source, ref: Yes




Outcome: Ever Stunted
Exposure:
Sex, ref: Female




Outcome: Ever Stunted
Exposure:
Small-for-gestational age, ref: Not SGA





Outcome: Ever Stunted
Exposure:
Single parent, ref: 0




Outcome: Wasted
Exposure:
Birth length (cm), ref: ≥ 50 cm










Outcome: Wasted
Exposure:
Breastfed hour after birth, ref: Yes





Outcome: Wasted
Exposure:
Father's age, ref: >=35












Outcome: Wasted
Exposure:
Improved floor, ref: Yes




Outcome: Wasted
Exposure:
Improved sanitation, ref: Yes







Outcome: Wasted
Exposure:
Mother's age, ref: [20-30)





Outcome: Wasted
Exposure:
Mother's BMI, ref: >=20 kg/m²












Outcome: Wasted
Exposure:
Birth order, ref: 1




Outcome: Wasted
Exposure:
Diarrhea <6 mo. (% days), ref: [0%, 2%]




Outcome: Wasted
Exposure:
Excl/Pred breastfed <6mo., ref: Yes




Outcome: Wasted
Exposure:
Rain quartile, ref: Opposite max rain




Outcome: Wasted
Exposure:
Sex, ref: Female




Outcome: Wasted
Exposure:
Small-for-gestational age, ref: Not SGA





Outcome: Wasted
Exposure:
Single parent, ref: 0




Outcome: Ever Wasted
Exposure:
Birth length (cm), ref: ≥ 50 cm







Outcome: Ever Wasted
Exposure:
Breastfed hour after birth, ref: Yes




Outcome: Ever Wasted
Exposure:
Father's age, ref: >=35







Outcome: Ever Wasted
Exposure:
Father's education, ref: High




Outcome: Ever Wasted
Exposure:
Father's height, ref: ≥ 162 cm












Outcome: Ever Wasted
Exposure:
Improved floor, ref: Yes




Outcome: Ever Wasted
Exposure:
Improved sanitation, ref: Yes










Outcome: Ever Wasted
Exposure:
Mother's weight, ref: ≥ 45 kg




Outcome: Ever Wasted
Exposure:
of children <5 in HH, ref: 1







Outcome: Ever Wasted
Exposure:
Birth order, ref: 1




Outcome: Ever Wasted
Exposure:
Safe water source, ref: Yes




Outcome: Ever Wasted
Exposure:
Sex, ref: Female




Outcome: Ever Wasted
Exposure:
Small-for-gestational age, ref: Not SGA




Outcome: Ever Wasted
Exposure:
Single parent, ref: 0




Outcome: Ever Wasted and Stunted
Exposure:
Birth length (cm), ref: ≥ 50 cm




Outcome: Ever Wasted and Stunted
Exposure:
Birthweight (kg), ref: ≥ 2500 g




Outcome: Ever Wasted and Stunted
Exposure:
Clean cooking fuel usage, ref: Yes




Outcome: Ever Wasted and Stunted
Exposure:
Breastfed hour after birth, ref: Yes




Outcome: Ever Wasted and Stunted
Exposure:
Father's age, ref: ≥ 35




Outcome: Ever Wasted and Stunted
Exposure:
Father's education, ref: High




Outcome: Ever Wasted and Stunted
Exposure:
Father's height, ref: ≥ 162 cm




Outcome: Ever Wasted and Stunted
Exposure:
Gestational age at birth, ref: Full or late term




Outcome: Ever Wasted and Stunted
Exposure:
Child delivered at home, ref: No




Outcome: Ever Wasted and Stunted
Exposure:
HH food security, ref: Food Secure




Outcome: Ever Wasted and Stunted
Exposure:
HH wealth, ref: Q4




Outcome: Ever Wasted and Stunted
Exposure: Improved floor, ref: Yes




Outcome: Ever Wasted and Stunted
Exposure: Improved sanitation, ref: Yes




Outcome: Ever Wasted and Stunted
Exposure:
Mother's age, ref: [20-30)




Outcome: Ever Wasted and Stunted
Exposure:
Mother's BMI, ref: ≥20 kg/m²




Outcome: Ever Wasted and Stunted
Exposure:
Mother's education, ref: High




Outcome: Ever Wasted and Stunted
Exposure:
Mother's height, ref: ≥ 150 cm



Outcome: Ever Wasted and Stunted
Exposure:
Mother's weight, ref: ≥ 45 kg




Outcome: Ever Wasted and Stunted
Exposure:
of children <5 in HH, ref: 1



Outcome: Ever Wasted and Stunted

Exposure:


of people in HH, ref: 3 or less




Outcome: Ever Wasted and Stunted

Exposure:


of rooms in HH, ref: 4+




Outcome: Ever Wasted and Stunted
Exposure:
Birth order, ref: 1




Outcome: Ever Wasted and Stunted
Exposure:
Safe water source, ref: Yes




Outcome: Ever Wasted and Stunted
Exposure:
Sex, ref: Female




Outcome: Ever Wasted and Stunted
Exposure:
Small-for-gestational age, ref: Not SGA




Outcome: Ever Wasted and Stunted
Exposure: Single parent, ref: 0




Outcome: Persistently Wasted
Exposure: Birth length (cm), ref: ≥ 50 cm




Outcome: Persistently Wasted
Exposure:
Birthweight (kg), ref: ≥ 2500 g




Outcome: Persistently Wasted
Exposure:
Clean cooking fuel usage, ref: Yes




Outcome: Persistently Wasted
Exposure:
Breastfed hour after birth, ref: Yes




Outcome: Persistently Wasted
Exposure:
Father's age, ref: >=35




Outcome: Persistently Wasted
Exposure:
Father's education, ref: High




Outcome: Persistently Wasted
Exposure:
Father's height, ref: ≥ 162 cm




Outcome: Persistently Wasted
Exposure:
Gestational age at birth, ref: Full or late term




Outcome: Persistently Wasted
Exposure:
Child delivered at home, ref: No




Outcome: Persistently Wasted
Exposure:
HH food security, ref: Food Secure




Outcome: Persistently Wasted
Exposure:
HH wealth, ref: Q4




Outcome: Persistently Wasted
Exposure: Improved floor, ref: Yes




Outcome: Persistently Wasted
Exposure: Improved sanitation, ref: Yes




Outcome: Persistently Wasted
Exposure:
Mother's age, ref: [20-30)




Outcome: Persistently Wasted
Exposure:
Mother's BMI, ref: ≥20 kg/m²




Outcome: Persistently Wasted
Exposure:
Mother's education, ref: High




Outcome: Persistently Wasted
Exposure:
Mother's height, ref: ≥ 150 cm



Outcome: Persistently Wasted
Exposure:
Mother's weight, ref: ≥ 45 kg




Outcome: Persistently Wasted
Exposure:
of children <5 in HH, ref: 1



Outcome: Persistently Wasted

Exposure:


of people in HH, ref: 3 or less




Outcome: Persistently Wasted

Exposure:


of rooms in HH, ref: 4+




Outcome: Persistently Wasted
Exposure:
Birth order, ref: 1




Outcome: Persistently Wasted
Exposure:
Sex, ref: Female




Outcome: Persistently Wasted
Exposure:
Small-for-gestational age, ref: Not SGA




Outcome: Persistently Wasted
Exposure:
Single parent, ref: 0




Outcome: Stunting recovery
Exposure:
Birth length (cm), ref: ≥ 50 cm
0-24 months




Outcome: Stunting recovery
Exposure:
Father's education, ref: High
0-24 months




Outcome: Stunting recovery
Exposure:
Gestational age at birth, ref: Full or late term
0-24 months




Outcome: Stunting recovery
Exposure:
HH wealth, ref: Q4
0-24 months



Outcome: Stunting recovery
Exposure:
Mother's age, ref: [20-30)
0-24 months



Outcome: Stunting recovery
Exposure:
Mother's education, ref: High
0-24 months




Outcome: Stunting recovery

Exposure:

Birth order, ref: 1

0-24 months



Outcome: Wasting recovery


Exposure:

Birth length (cm), ref: ≥ 50 cm


0-6 months

6-24 months


0-24 months




Outcome: Wasting recovery
Exposure:
Birthweight (kg), ref: ≥ 2500 g




Outcome: Wasting recovery
Exposure:
Clean cooking fuel usage, ref: Yes







Outcome: Wasting recovery
Exposure:
Breastfed hour after birth, ref: Yes





Outcome: Wasting recovery
Exposure:
Father's age, ref: >=35












Outcome: Wasting recovery
Exposure:
Improved floor, ref: Yes




Outcome: Wasting recovery
Exposure:
Improved sanitation, ref: Yes





Outcome: Wasting recovery
Exposure:
Mother's age, ref: [20-30)




Outcome: Wasting recovery
Exposure:
Mother's BMI, ref: ≥ 20 kg/m²






Outcome: Wasting recovery
Exposure:
Mother's weight, ref: ≥ 45 kg




Outcome: Wasting recovery
Exposure:
of children <5 in HH, ref: 1



Outcome: Wasting recovery

Exposure:


of people in HH, ref: 3 or less




Outcome: Wasting recovery

Exposure:


of rooms in HH, ref: 4+




Outcome: Wasting recovery
Exposure:
Birth order, ref: 1



Outcome: Wasting recovery
Exposure:
Sex, ref: Female



Outcome: Wasting recovery
Exposure:
Small-for-gestational age, ref: Not SGA



Outcome: Wasting recovery
Exposure:
Single parent, ref: 0

