

# Large-scale detection and characterization of interchromosomal rearrangements in normozoospermic bulls using massive genotype and phenotype datasets

by Jeanlin Jourdain *et al.*

## Table of contents:

### File Supplemental\_Figs.pdf:

Supplemental Figure 1. Information on t(5;28) and bull Fa	1-3
Supplemental Figure 2. Information on t(2;22) and bull Fa	4-6
Supplemental Figure 3. Information on t(10;12) and bull Im	7
Supplemental Figure 4. Information on t(1;11) and bull Ja	8-10
Supplemental Figure 5. Information on t(5;28) and bull Le	11-13
Supplemental Figure 6. Information on t(3;8) and bull Ma	14-18
Supplemental Figure 7. Information on inv ins (4;8) and bull Nt	19-24
Supplemental Figure 8. Information on t(12;23) and bull No	25
Supplemental Figure 9. Information on Rob(18;21) and bull Ob	26
Supplemental Figure 10. Information on t(24;29) -der29 and bull Ou	27-31
Supplemental Figure 11. Information on t(12;23) and bull Qu	32
Supplemental Figure 12. Information on bull Ra	33
Supplemental Figure 13. Information on t(2;13) and bull Sa	34
Supplemental Figure 14. Distribution of linkage disequilibrium in chromosome	35

### File Supplemental\_Tables1-4.pdf:

Supplemental Table 1. Raw technical numbers for affected bulls.	1
Supplemental Table 2. Number of eligible bull cohorts per breed.	1
Supplemental Table 3. Analysis of linkage disequilibrium around breakpoints.	2
Supplemental Table 4. Repeated regions around breakpoints	2

### File Supplemental\_Table5.xlsx:

Supplemental Table 5. List of genes potentially affected by breakpoints.
--