

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Long-Term Exposure to Walkable Residential Neighborhoods and Risk of Obesity-Related Cancer in the New York University Women's Health Study (NYUWHS)

Sandra India-Aldana, Andrew G. Rundle, James W. Quinn, Tess V. Clendenen, Yelena Afanasyeva, Karen L. Koenig, Mengling Liu, Kathryn M. Neckerman, Lorna E. Thorpe, Anne Zeleniuch-Jacquotte, and Yu Chen

Table of Contents

Table S1. First Incident Obesity-related Cancers in the NYUWHS (n=13,240).

Table S2. Linearity Test using Cubic splines for the Association between Neighborhood Walkability and Obesity-related Cancer in the NYUWHS (n=13,240).

Table S3. Testing Proportional Hazards (PH) Assumption in the Association between Neighborhood Walkability and Obesity-related Cancer in the NYUWHS (n=13,240).

Table S4. NYUWHS Demographic and Lifestyle Characteristics by History of Moving from Baseline Residence, N=13,240.

Table S5. Neighborhood Walkability Exposure Availability and Person-Time Years in the NYUWHS.

Table S6. Obesity-related Cancer Hazard Ratios based on Neighborhood Walkability Sensitivity Analyses.

Table S7. Obesity-related Cancer Hazard Ratios based on Neighborhood Walkability Sensitivity Analyses Using Different Time-Scales (n=13,240).

Table S8. Obesity-related Cancer Hazard Ratios based on Median and Continuous (Per IQR) Neighborhood Walkability (n=13,240).

Table S9. Stratified Analyses by Age, BMI, and Parity for Additional Frequent Cancer Subtypes (n=13,240).

Table S10. Stratified Analyses by Poverty, DASH Diet, and Smoking Status for Additional Frequent Cancer Subtypes (n=13,240).

Table S11. Stratified Analyses by Race for Additional Frequent Cancer Subtypes in the NYUWHS (n=11,608).

Table S12. Stratified Analyses by Education for Additional Frequent Cancer Subtypes in the NYUWHS (n=10,842).

Table S13. Stratified Analyses by Median Outdoor Walking at Baseline and History of Diabetes for Most Frequent Cancer Subtypes in the NYUWHS (n=13,240).

Table S14. Postmenopausal Breast Cancer Competing Risk Hazard Ratios based on Neighborhood Walkability by ER/PR Status.

Figure S1. Follow-up Questionnaires' Timeline in the NYUWHS.

Figure S2. Time-varying Exposure Characterization Scenario for NYUWHS Participant with Long Follow-Up.

References