

Supplementary Table 1. Sequences of primers used for qRT-PCR analysis.

	Forward (5'-3')	Reverse (5'-3')
<i>Actin</i>	GGCTGTATCCCCTCCATCG	CCAGTTGGTAACAATGCCATGT
<i>Ppara</i>	ACGGCAATGGCTTTATCA	CGCTGCGTCGGACTCGGT
<i>Fatp2</i>	TCCTCCAAGATGTGCGGTACT	TAGGTGAGCGTCTCGTCTCG
<i>Fatp5</i>	CTACGCTGGCTGCATATAGATG	CCACAAAGGTCTCTGGAGGAT
<i>Cpt1a</i>	ACCACTGGCCGCATGT	CTCCATGGCGTAGTAGTTGCT
<i>Cpt2</i>	CAGCACAGCATCGTACCCA	TCCAATGCCGTTCTCAAAT
<i>Cox4</i>	CGGCGTGACTACCCCTTG	TGAGGGATGGGGCCATACA
<i>Cyto C</i>	CCAAATCTCCACGGTCTGTTC	ATCAGGGTATCCTCTCCCCAG
<i>Acadl</i>	GCATCAACATCGCAGAGAAA	ACGCTTGCTCTTCCCAAGTA
<i>Acadm</i>	GCTAGTGGAGCACCAAGGAG	CCAGGCTGCTCTCTGGTAAC
<i>Pck1</i>	GAAGGACAAAGATGGCAAGTT	CGTTTTCTTAGGGATGTAGC
<i>G6pc</i>	CTGAGCGCGGGCATCATAAT	GATTCTTAGGATCGCCCAGAAAG
<i>Hmgcs2</i>	GAAGAGAGCGATGCAGGAAAC	GTCCACATATTGGGCTGGAAA
<i>Hmgcl</i>	CAGGTGAAGATCGTGGAAGTC	GGAGCCCTGCTTCGGAAC
<i>Bdh1</i>	ACAAGACACACGCTGTTGTTT	CTCTTCAAGCTGTCCAGTTCC
<i>Gck</i>	GTCTTTTGCAACACTCAGCCA	ATCCGGCTCATCACCTTCTTC
<i>Pfkf</i>	GGAGGCGAGAACATCAAGCC	CGGCCTTCCCTCGTAGTGA
<i>Pklr</i>	TGGCATCGAAAGTGGAAGC	GATGTGGGACTATGGGAGGG
<i>Srebp1c</i>	GGAGCCATGGATTGCACATT	GCTTCCAGAGAGGAGGCCAG
<i>Acc1</i>	AGCTGATCCTGCGAACCT	GCCAAGCGGATGTAAACT
<i>Fasn</i>	TCCAAGACTGACTCGGCTACTGAC	GCAGCCAGGTTCGGAATGCTATC
<i>Elovl6</i>	AAGCAGTTCAACGAGAACGAA	CGTACAGCGCAGAAAACAGG
<i>Scd1</i>	TTCTTGCGATACTCTGGTGC	CGGGATTGAATGTTCTTGTCTG
<i>Hmgcs1</i>	AATTGGGCCAAACGCTCCT	TCTCGTACAAGAGAACTGGCTA
<i>Hmgcr</i>	CTGGAATTATGAGTGCCCCAAA	ACGACTGTACTGAAGACAAAGC
<i>Fdft1</i>	ATGGAGTTCGTCAAGTGCTAGG	CGTGCCGTATGTCCCATC
<i>Lss</i>	TCGTGGGGGACCCTATAAAAC	CGTCCTCCGCTTGATAATAAGTC