

**S1 Table.** Univariable and multivariable analyses for the risk factors for overall survival in patients with ypT+NO patients

	Univariable analysis		Multivariable analysis	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Male sex	0.734 (0.260-2.070)	0.560	0.825 (0.242-2.811)	0.759
Age	0.990 (0.957-1.025)	0.591	0.980 (0.940-1.022)	0.354
ypStage (vs. II)				
III	1.196 (0.621-2.300)	0.591	1.119 (0.557-2.247)	0.752
IV	2.593 (0.734-9.150)	0.139	1.774 (0.431-7.296)	0.427
FEV <sub>1</sub>	0.682 (0.349-1.331)	0.262	-	-
DLC <sub>o</sub>	1.006 (0.986-1.027)	0.539	-	-
ASA (vs. I)				
II	2.948 (1.186-7.326)	0.020	3.023 (1.164-7.851)	0.023
III	8.782 (2.574-29.968)	0.001	11.399 (3.008-43.197)	< 0.001
Adjuvant chemotherapy	1.218 (0.342-4.342)	0.760	-	-

ASA, American Society of Anesthesiology; CI, confidence interval; DLC<sub>o</sub>, diffusing capacity of the lungs for carbon monoxide; FEV<sub>1</sub>, forced expiratory volume in 1 second; HR, hazard ratio.