

S2 Table. Univariable and multivariable analyses for the risk factors for overall survival in patients with ypT0N+ patients

	Univariable analysis		Multivariable analysis	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Male sex	0.549 (0.167-1.805)	0.323	0.425 (0.095-1.894)	0.262
Age	0.992 (0.950-1.035)	0.723	1.000 (0.950-1.053)	0.977
ypStage (vs. II)				
III	1.334 (0.612-2.904)	0.467	1.764 (0.637-4.883)	0.275
IV	10.187 (1.244-83.381)	0.030	3.794 (0.317-45.357)	0.292
FEV ₁	1.111 (0.615-2.006)	0.727	-	-
DLC _o	0.983 (0.963-1.003)	0.099	0.982 (0.958-1.005)	0.138
ASA (vs. I)				
II	0.968 (0.460-2.038)	0.934	0.943 (0.180-4.928)	0.945
III	2.811 (0.933-8.472)	0.066	1.406 (0.211-9.358)	0.724
Adjuvant chemotherapy	1.027 (0.531-1.986)	0.936	-	-

ASA, American Society of Anesthesiology; CI, confidence interval; DLC_o, diffusing capacity of the lungs for carbon monoxide; FEV₁, forced expiratory volume in 1 second; HR, hazard ratio.