

S2 Table. Absolute lymphocyte count (ALC) and changes in study population during study

Patient	Baseline	ALC ($\times 10^3/\mu\text{L}$)	ALC ($\times 10^3/\mu\text{L}$)
	ALC ($\times 10^3/\mu\text{L}$) (before cycle 1)	after proton treatment 1 (before cycle 2)	after proton treatment 2 (before cycle 3)
1	1.42	0.81	no data
2	1.64	1.89	2.13
3	1.79	1.33	1.35
4	1.32	0.98	1.34
5	0.67	3.20	0.61
6	1.36	1.43	1.33
7	1.21	0.78	1.10
8	1.10	0.61	0.81
9	1.56	1.09	0.94
10	0.53	0.97	0.52
11	1.35	0.76	No data
12	0.52	0.76	0.56
13	1.87	2.00	2.19
14	0.90	0.59	0.60
15	0.75	0.77	No data
16	1.25	0.96	1.20
17	1.51	1.15	0.87
18	0.90	0.88	0.46
19	2.13	2.08	1.68
20	0.50	0.60	0.63
21	0.82	0.76	0.31
22	1.32	1.27	1.26
23	0.85	0.85	0.96
24	1.00	0.66	No data
25	1.53	0.59	0.18