

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Prenatal Exposure to Multiple Endocrine-Disrupting Chemicals and Childhood BMI Trajectories in the INMA Cohort Study

Parisa Montazeri, Nuria Güil-Oumrait, Sandra Marquez, Lourdes Cirugeda, Andrea Beneito, Mònica Guxens, Aitana Lertxundi, Maria-Jose Lopez-Espinosa, Loreto Santa-Marina, Jordi Sunyer, Maribel Casas, and Martine Vrijheid

Table of Contents

Table S1. Details regarding laboratories and LOD or LOQ by sampling period for Gipuzkoa subcohort of the INMA birth cohort study.

Table S2. Details regarding laboratories and LOD or LOQ by sampling period for Sabadell subcohort of the INMA birth cohort study.

Table S3. Details regarding laboratories and LOD or LOQ by sampling period for Valencia subcohort of the INMA birth cohort study.

Table S4. Description of the growth trajectory modeling used in the INMA birth cohort study.

Table S5. Characteristics of the study population, unimputed and imputed data for the INMA birth cohort study.

Table S6. Associations between EDCs and childhood BMI trajectories from single exposure models using multinomial regression models and expressed as relative risk ratios, for the INMA birth cohort study.

Table S7. Associations between EDCs and childhood BMI trajectories from single exposure models using multinomial regression models and expressed as relative risk ratios for the INMA birth cohort study, complete case analysis.

Table S8. Associations between EDCs and childhood BMI trajectories from single exposure models using multinomial regression models and expressed as relative risk ratios for the INMA birth cohort study, by sex [females (n = 927) / males (n = 984)].

Table S9. Associations between EDCs and childhood BMI growth trajectory classes from single exposure models for the INMA birth cohort study by socioeconomic status, high (n = 613) / middle (n = 497) / low (n = 777).

Table S10. Associations between EDC mixture and childhood BMI trajectories from mixture models using BWQS regression (n = 1,911), for the INMA birth cohort study.

Table S11. Class-specific mean predicted trajectory values and their confidence intervals for each BMI trajectory class depicted in Figure 1, by select age groups.

Figure S1. Directed acyclic graph of the association between prenatal exposure to EDCs and zBMI growth trajectories.

Figure S2-3. Pairwise Correlations of EDC exposures for the INMA birth cohort study.