

Supplemental Table 1 : Original CoMiSS with BITSS

| Crying | |
|--------------------|---|
| ≤1 hour/day | 0 |
| 1 to 1.5 hours/day | 1 |
| 1.5 to 2 hours/day | 2 |
| 2 to 3 hours/day | 3 |
| 3 to 4 hours/day | 4 |
| 4 to 5 hours/day | 5 |
| ≥5 hours/day | 6 |

| Regurgitation | |
|---|---|
| 0 to 2 episodes/day | 0 |
| ≥3 to ≤5 of small volume | 1 |
| >5 episodes of >1 coffee spoon | 2 |
| >5 episodes of ± half of the feed in <half of the feedings | 3 |
| Continuous regurgitations of small volume >30min after each feeding | 4 |
| Regurgitation of half to complete volume of a feeding in at last half of the feedings | 5 |
| Regurgitation of the "complete feeding" after each feeding | 6 |

| Stools (Bristol scale) | |
|--|---|
| Type 1 and 2 (hard stools) | 4 |
| Type 3 and 4 (normal stools) | 0 |
| Type 5 (soft stool) | 2 |
| Type 6 (liquid stool, if unrelated to infection) | 4 |
| Type 7 (watery stools) | 6 |

| Skin Symptoms | | | | |
|----------------------------|--------|------|----------|--------|
| Atopic eczema | absent | mild | moderate | severe |
| Head, neck and trunk | 0 | 1 | 2 | 3 |
| Arms, hands, legs and feet | 0 | 1 | 2 | 3 |

| Urticaria | no | yes |
|-----------|----|-----|
| | 0 | 6 |

| Respiratory symptoms | |
|-----------------------------|---|
| No respiratory symptoms | 0 |
| Slight symptoms | 1 |
| Mild symptoms | 2 |
| Severe symptoms | 3 |