



Supplemental Figure 1 The flow chart

IgAN-NS, the IgAN with NS; IgAN-NR, the IgAN with nephrotic-range proteinuria; MCD, minimal change disease; MN, Membranous nephropathy.

Table S1 The percentage of missing data of 222 cases in this study

Factors	The percentage of missing data
Age,y	0%
Male	0%
Follow-up duration,m	0%
eGFR	0%
Body mass index	2.0%
Systolic blood pressure, mmHg	6.0%
Diastolic blood pressure, mmHg	6.0%
Mean arterial pressure, mmHg	6.0%
Hemoglobin, g/L	2.7%
Serum creatinine, mg/dL	0%
Serum albumin, g/L	0%
Total cholesterol, mmol/L	1.4%
Triglyceride, mmol/L	1.4%
HDL-cholesterol, mmol/L	1.4%
LDL-cholesterol, mmol/L	1.4%
Uric acid,mmol/L	8.1%
Fibrinogen, g/L	2.3%
C-reactive protein, g/L	9.9%
Serum IgA, g/L	1.4%
Proteinuria, g/24h	0%
Urine red blood cell count, RBC/ μ L	0%
Urine white blood cell count, WBC/ μ L	0%
Therapy	
Renin-angiotensinsystem inhibito	0%
Steroid	0%
Immunosuppressant	0%