

Table S2. Diagnostic accuracy of serum AMA for individual study.

Author	Year	TP	FP	FN	TN	Sensitivity	Specificity
Assassi ^[1]	2009	7	9	9	792	43.8%	98.9%
Bandin ^[2]	1996	270	5	15	492	94.7%	99.0%
Bargou ^[3]	2008	54	0	1	51	98.2%	100.0%
Cavazzana ^[4]	2011	14	5	9	173	60.9%	97.2%
Chung ^[5]	2016	8	2	0	24	80.0%	100.0%
Dighiero ^[6]	1987	21	18	6	666	77.8%	97.4%
Gabeta ^[7]	2007	76	0	27	111	73.8%	100.0%
Guatibonza-García ^[8]	2021	19	10	24	32	44.2%	76.2%
Han ^[8]	2017	28	9	7	99	91.4%	87.0%
Hu ^[9]	2011	183	10	15	75	92.4%	88.2%
Jong-Hon ^[10]	2001	49	0	14	291	77.8%	100.0%
Lee ^[11]	1983	9	2	1	10	90.0%	83.3%
Liu and Liu ^[12]	2010	4	31	0	8901	100.0%	99.7%
Liu and Norman ^[13]	2010	985	65	190	1167	83.8%	94.7%
Muratori ^[14]	2004	91	7	36	259	71.7%	97.4%
Nagai ^[15]	1983	16	7	0	97	100.0%	93.3%
Oertelt ^[16]	2007	96	0	24	124	80.0%	100.0%
Romero-Gómez ^[17]	2004	95	4	41	34	69.9%	89.5%
Sakugawa ^[18]	2003	6	4	1	111	85.7%	96.5%
Sun ^[19]	2019	72	0	55	100	56.7%	100.0%
Villalta ^[20]	2015	50	8	187	4	86.2%	97.9%
Wang ^[21]	2015	58	0	0	30	100.0%	100.0%
Yang J ^[22]	2016	98	4	18	120	84.5%	96.8%
Yang Z ^[23]	2012	102	13	13	787	88.7%	98.4%

TP =True positive, FP = False positive, FN = False negative, TN = True negative.