

Table S3. Diagnostic accuracy of serum M2 subtype for individual study.

Author	Year	TP	FP	FN	TN	Sensitivity	Specificity
Assassi ^[1]	2009	13	43	3	758	81.3%	94.6%
Bargou ^[3]	2008	51	4	4	47	92.7%	92.2%
Dighiero ^[6]	1987	27	38	0	646	100.0%	94.4%
Gabeta ^[7]	2007	84	0	19	111	81.6%	100.0%
Han ^[8]	2017	32	9	3	99	91.4%	91.7%
Hu ^[9]	2011	173	20	25	65	87.4%	76.5%
Imura-Kumada ^[24]	2012	22	15	0	188	100.0%	92.6%
Lu ^[25]	2017	33	0	16	60	67.3%	100.0%
Milkiewicz ^[26]	2009	165	14	49	53	77.1%	79.1%
Muratorj ^[14]	2004	100	6	27	260	78.7%	97.7%
Nagai ^[15]	1983	16	2	0	102	100.0%	98.1%
Sun ^[19]	2019	66	0	61	100	52.0%	100.0%
Tang ^[27]	2017	60	0	0	37	100.0%	100.0%
Villalta ^[20]	2015	45	13	187	4	77.6%	97.9%
Wang ^[21]	2015	55	3	3	27	94.8%	90.0%
Yang J ^[22]	2016	98	3	18	121	84.5%	97.6%

TP =True positive, FP = False positive, FN = False negative, TN = True negative