

B


C



Name and Proportion of the Cellular Component (Inner Ring)



Name and Proportion of the Molecular Function (Inner Ring)



- GO1 voltage-gated calcium channel activity, 26.9%
- GO2 cell adhesion molecule binding, 8.7%
- GO3 olfactory receptor activity, 6.8%
- GO4 spectrin binding, 5.2%
- GO5 carbohydrate binding, 4.5%
- GO6 G-rich strand telomeric DNA binding, 4.1%
- GO7 single-stranded DNA binding, 4.1%
- GO8 5'-3' DNA helicase activity, 3.9%
- GO9 glucan 1,3-alpha-glucosidase activity, 3.8%
- GO10 telomeric repeat-containing RNA binding, 3.0%
- single-stranded RNA binding, 2.9%
- methyl-CpG binding, 2.6%
- tRNA (guanine-N2-)-methyltransferase activity, 2.3%
- S100 protein binding, 2.1%
- nitric-oxide synthase activity, 2.1%
- latrotoxin receptor activity, 2.1%
- tRNA (guanine-N3-)-methyltransferase activity, 1.9%
- odorant binding, 1.9%
- 4 iron, 4 sulfur cluster binding, 1.8%
- ammonia-lyase activity, 1.7%
- tRNA (guanine-N4-)-methyltransferase activity, 1.7%
- 3',5'-cyclic-AMP phosphodiesterase activity, 1.6%
- tRNA (guanine-N5-)-methyltransferase activity, 1.6%
 - damaged DNA binding, 1.3%
 - NADP binding, 1.3%