

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

E-reference 1: Distribution of baseline rectal swab and first-void urine *Chlamydia trachomatis* transcription-mediated amplification results, with stratification by six- and twelve-month follow-up encounter testing results, within the RADAR longitudinal cohort investigation.

	Frequency of <i>C. trachomatis</i> Screening Result					
Baseline Rectal Swab Screen	<i>Six-month Follow-up (rectal swab)</i>			<i>Twelve-month Follow-up (rectal swab)</i>		
	Non-reactive	Reactive	Total	Non-reactive	Reactive	Total
Non-reactive	652	46	698	572	43	615
Reactive	53	13	66	45	7	52
Total	705	59	764	617	50	667
Baseline Urine Screen	<i>Six-month Follow-up (urine)</i>			<i>Twelve-month Follow-up (urine)</i>		
	Non-reactive	Reactive	Total	Non-reactive	Reactive	Total
Non-reactive	755	6	761	655	9	664
Reactive	8	2	10	9	0	9
Total	763	8	771	664	9	673

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

E-reference 2: Distribution of baseline rectal swab and first-void urine *Neisseria gonorrhoeae*
transcription-mediated amplification results, with stratification by six- and
twelve-month follow-up encounter testing results, within the RADAR
longitudinal cohort investigation.

	Frequency of <i>N. gonorrhoeae</i> Screening Result					
Baseline Rectal Swab Screen	<i>Six-month Follow-up (rectal swab)</i>			<i>Twelve-month Follow-up (rectal swab)</i>		
	Non-reactive	Reactive	Total	Non-reactive	Reactive	Total
Non-reactive	674	35	709	593	22	615
Reactive	44	11	55	43	11	54
Total	718	46	764	636	33	669
Baseline Urine Screen	<i>Six-month Follow-up (urine)</i>			<i>Twelve-month Follow-up (urine)</i>		
	Non-reactive	Reactive	Total	Non-reactive	Reactive	Total
Non-reactive	761	3	764	664	4	668
Reactive	6	1	7	5	0	5
Total	767	4	771	669	4	673

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

E-reference 3: Distribution of baseline rectal swab and first-void urine *Trichomonas vaginalis*
transcription-mediated amplification results, with stratification by six- and
twelve-month follow-up encounter testing results, within the RADAR
longitudinal cohort investigation.

	Frequency of <i>T. vaginalis</i> Screening Result					
Baseline Rectal Swab Screen	<i>Six-month Follow-up (rectal swab)</i>			<i>Twelve-month Follow-up (rectal swab)</i>		
	Non-reactive	Reactive	Total	Non-reactive	Reactive	Total
Non-reactive	754	8	762	659	9	668
Reactive	5	1	6	2	2	4
Total	759	9	768	661	11	672
Baseline Urine Screen	<i>Six-month Follow-up (urine)</i>			<i>Twelve-month Follow-up (urine)</i>		
	Non-reactive	Reactive	Total	Non-reactive	Reactive	Total
Non-reactive	769	1	770	672	0	672
Reactive	0	1	1	0	1	1
Total	769	2	771	672	1	673

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

E-reference 4: Distribution of baseline rectal swab and first-void urine *Mycoplasma genitalium* transcription-mediated amplification results, with stratification by six- and twelve-month follow-up encounter testing results, within the RADAR longitudinal cohort investigation.

	Frequency of <i>M. genitalium</i> Screening Result					
Baseline Rectal Swab Screen	<i>Six-month Follow-up (rectal swab)</i>			<i>Twelve-month Follow-up (rectal swab)</i>		
	Non-reactive	Reactive	Total	Non-reactive	Reactive	Total
Non-reactive	535	66	601	463	69	532
Reactive	64	103	167	78	62	140
Total	599	169	768	541	131	672
Baseline Urine Screen	<i>Six-month Follow-up (urine)</i>			<i>Twelve-month Follow-up (urine)</i>		
	Non-reactive	Reactive	Total	Non-reactive	Reactive	Total
Non-reactive	683	19	702	594	20	614
Reactive	15	54	69	17	40	57
Total	698	73	771	611	60	671