

Supplementary data

CNS drugs

Long term treatment over 52 weeks with monthly fremanezumab in drug resistant migraine: a prospective, multicenter study

Valeria Caponnetto¹, Antonio Russo², Marcello Silvestro², Alessandro Tessitore², Roberto De Icco^{3,4}, Gloria Vaghi^{3,4}, Grazia Sances⁴, Cristina Tassorelli^{3,4}, Carlo Baraldi⁵, Flavia Lo Castro⁵, Simona Guerzoni⁵, Maria Pia Prudenzano⁶, Adriana Fallacara⁶, Martino Gentile⁶, Raffaele Ornello¹, Agnese Onofri¹, Andrea Burgalassi⁷, Alberto Chiarugi^{7,8}, Francesco De Cesaris⁷, Antonio Granato⁹, Alfonsina Casalena¹⁰, Marina De Tommaso¹¹, Edoardo Mampreso¹², Paola Merlo¹³, Gianluca Coppola¹⁴, Stefania Battistini¹⁵, Valentina Rebecchi¹⁶, Innocenzo Rainero¹⁷, Federica Nicoletta Sepe¹⁸, Giorgio Dalla Volta¹⁹, Simona Sacco¹, Pierangelo Geppetti^{7,8}, Luigi Francesco Iannone^{7*}, on behalf of the Italian Headache Registry (RICE) Study Group.

¹Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, L'Aquila, Italy.

²Headache Centre, Department of Advanced Medical and Surgical Sciences, University of Campania "Luigi Vanvitelli", Naples, Italy.

³Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy.

⁴Headache Science & Neurorehabilitation Center, IRCCS Mondino Foundation, Pavia, Italy.

⁵Department of Specialist Medicines, Digital and Predictive Medicine, Pharmacology and Clinical Metabolic Toxicology-Headache Center and Drug Abuse, Laboratory of Clinical Pharmacology and Pharmacogenomics, AOU Policlinico di Modena, Modena, Italy.

⁶Headache Center, Department of Basic Medical Sciences, Neurosciences and Sense Organs, University of Bari, Bari, Italy.

⁷Headache Center and Clinical Pharmacology Unit, Careggi University Hospital, Florence, Italy.

⁸Section of Clinical Pharmacology and Oncology, Department of Health Sciences, University of Florence, Florence, Italy.

⁹Headache Centre, Clinical Unit of Neurology, School of Neurology, Department of Medicine, Surgery and Health Sciences, University Hospital and Health Services of Trieste, ASUGI, University of Trieste, Italy.

¹⁰Neurology Unit, "G. Mazzini" Hospital, Teramo, Italy.

¹¹Department of Basic Medical Science, Neuroscience and Sense Organs, University of Bari Aldo Moro, Bari, Italy.

¹²Headache Centre, Neurology - Euganea, Padua Health Unit, Padua, Italy.

¹³Neurological Unit, Headache Centre, Humanitas Gavazzeni, Bergamo, Italy.

¹⁴Department of Medico-Surgical Sciences and Biotechnologies, Sapienza University of Rome Polo Pontino ICOT, Latina, Italy.

¹⁵ Headache Center, Neurology and Clinical Neurophysiology Unit, Siena University Hospital Department of Medical, Surgical and Neurological Sciences, University of Siena, Siena, Italy.

¹⁶ Neurology and Stroke Unit, ASST Sette Laghi, Varese, Italy.

¹⁷Headache Center, Department of Neuroscience "Rita Levi Montalcini", University of Torino, Italy.

¹⁸ Neurology Department, Azienda Ospedaliera Nazionale Santi Antonio e Biagio e Cesare Arrigo, Alessandria, Italy.

¹⁹ U.O. Neurologia, Headache Center Istituto Clinico "Città di Brescia", Gruppo San Donato Brescia, Italy.

*Author for correspondence:

Luigi Francesco Iannone

Department of Health Sciences

University of Florence, Viale Pieraccini 6, Florence, Italy

Luigifrancesco.iannone@unifi.it

Table S1. Concomitant and prior preventive treatments.

	Overall population (n =83)	HFEM (n=16)	CM (n=67)
Patients with concomitant preventive treatment, n (%)	27 (32.5)	4 (25.0)	23 (34.3)
Concomitant preventive treatment, mean \pm SD	1.3 \pm 0.6	1.5 \pm 0.5	1.2 \pm 0.6
Prior preventive classes failures, mean \pm SD	4.0 \pm 1.4	3.5 \pm 1.1	4.1 \pm 1.5
Prior preventive classes non-tolerated, mean \pm SD	0.3 \pm 0.6	0.3 \pm 0.7	0.2 \pm 0.5
Prior preventive class failures			
<3 classes	12 (14.4)	3 (18.8)	9 (13.5)
\geq 3 classes	71 (85.6)	13 (81.5)	58 (86.5)
\geq 4 classes	54 (69.6)	8 (50.0)	41 (61.1)
\geq 5 classes	32 (38.6)	3 (18.7)	29 (43.2)
\geq 6 classes	13 (15.7)	1 (6.2)	12 (17.9)
\geq 7 classes	2 (2.4)	0	2 (2.9)
Drug Classes			
Beta-blockers	64 (78.1)	11 (68.8)	53 (80.3)
Tricyclic antidepressant	74 (89.2)	15 (93.8)	59 (88.1)
Calcium channel blockers	53 (63.9)	13 (81.3)	40 (59.7)
Antiseizure medications	65 (78.3)	11 (68.8)	54 (80.6)
SSRIs/SNRIs	15 (18.1)/ 8 (9.6)	3 (18.8)/ 1 (6.3)	12 (17.9)/ 7 (10.4)
OnabotulinumtoxinA	35 (42.2)	2 (12.5)	33 (49.3)
Angiotensin receptor antagonists	18 (21.7)	1 (6.3)	17 (25.4)

Abbreviations: CM, chronic migraine; HFEM, high frequency episodic migraine; SD, standard deviation; SSRI, selective serotonin reuptake inhibitors; SNRI, serotonin-norepinephrine reuptake inhibitor. Percentages are expressed on column total.

Table S2. Adverse events reported through 12 months of follow-up.

	Overall population (n =83)	HFEM (n=16)	CM (n=67)
Patients reporting at least one AE, n (%)	8 (9.6)	2 (12.5)	6 (8.9)
Overall AEs rate, n (%)	10 (12.0)	3 (18.7)	7 (10.4)
AEs leading to drug discontinuation, n (%)	1 (1.2)	1 (6.2)	-
AEs reported			
Constipation, n (%)	6 (7.2)	2 (12.5)	4 (5.9)
Injection site reaction, n (%)	3 (3.6)	1 (6.2) ^a	2 (2.9)
Effluvium, n (%)	1 (1.2)	-	1 (1.4)

Abbreviations: AE, adverse event; CM, chronic migraine; HFEM, high frequency episodic migraine. Percentages are expressed on column total. HFEM and CM are part of the overall population. ^aleading to discontinuation.

Table S3. Changes in migraine-related variables during fremanezumab treatment in the overall, HFEM and CM populations (LOCF).

	Median (IQR)		Median difference (IQR)	
	Baseline	3 months	6 months	12 months
Overall population (n=83)				
Monthly headache days	20.0 (15.0)	-9.2 (12.0)	-9.0 (14.0)	-9.0 (16.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
Analgesic medication days	15.0 (18.0)	-9.0 (11.0)	-9.0 (12.0)	-9.0 (13.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
Analgesic medication number	20.0 (20.0)	-10.0 (18.0)	-11.0 (12.0)	-12.0 (17.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
MIDAS score	64.0 (60.0)	-35.0 (49.0)	-35.0 (54.0)	-35.0 (54.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
HIT-6 score	67.0 (7.0)	-10.0 (15.0)	-12.0 (16.0)	-12.0 (17.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
HFEM (n=16)				
Monthly headache days	10.0 (2.0)	-5.0 (5.2)	-6.0 (4.0)	-6.0 (5.2)
<i>p value</i>	-	0.001	0.001	0.001
Analgesic medication days	9.0 (2.0)	-7.0 (6.0)	-6.5 (6.5)	-5.0 (6.0)
<i>p value</i>	-	0.005	0.004	0.004
Analgesic medication number	10.0 (2.0)	-6.0 (7.5)	-7.5 (8.0)	-6.0 (7.5)
<i>p value</i>	-	0.007	0.005	0.006
MIDAS score	30.5 (31)	-18.0 (20.5)	-18.0 (35.2)	-18.0 (29.7)
<i>p value</i>	-	0.003	0.002	0.002
HIT-6 score	67.0 (6.0)	-10.0 (13.2)	-20.0 (16.0)	-15.0 (19.2)
<i>p value</i>	-	0.001	<0.001	<0.001
CM (n=67)				
Monthly headache days	25.0 (14.0)	-11.0 (14.0)	-11.0 (17.0)	-11.0 (18.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
Analgesic medication days	20.0 (18.0)	-10.0 (12.0)	-10.0 (13.0)	-10.0 (14.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
Analgesic medication number	25.0 (18.0)	-13.0 (21.0)	-13.0 (14.0)	-14.0 (17.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
MIDAS score	80.0 (65.0)	-47.0 (60.0)	-45.0 (57.0)	-45.0 (53.0)
<i>p value</i>	-	<0.001	<0.001	<0.001
HIT-6 score	67.0 (7.0)	-10.0 (15.0)	-11.0 (18.0)	-12.0 (18.0)
<i>p value</i>	-	<0.001	<0.001	<0.001

Abbreviations: CM, chronic migraine; HFEM, high frequency episodic migraine; IQR, interquartile range; LOCF, last observation carried forward; HIT-6, Headache Impact Test 6; MIDAS, Migraine Disability Assessment; SD, standard deviation. Baseline values are the same of observed data analysis but reported for data consistency. The changes are calculated using the Wilcoxon signed-rank test and expressed as median reduction [IQR]. All *p* values and reductions are vs baseline. Significant *p* values are reported in bold. IQR, interquartile range.

Table S4. Response rate during fremanezumab treatment in HFEM, CM and overall populations (LOCF).

	3 months n (%)	6 months n (%)	12 months n (%)
Overall population (n=83)			
Response rate $\geq 30\%$	57 (68.7)	60 (72.3)	58 (69.9)
Response rate $\geq 50\%$	42 (50.6)	51 (61.4)	53 (63.8)
Response rate $\geq 75\%$	19 (22.9)	26 (31.3)	25 (30.1)
Response rate 100%	2 (2.4)	4 (4.8)	5 (6.0)
HFEM (n=16)			
Response rate $\geq 30\%$	11 (68.7)	13 (81.2)	12 (75.0)
Response rate $\geq 50\%$	9 (56.3)	10 (62.5)	11 (68.7)
Response rate $\geq 75\%$	2 (25.0)	6 (37.5)	5 (31.3)
Response rate 100%	2 (12.5)	2 (12.5)	2 (12.5)
CM (n=67)			
Response rate $\geq 30\%$	46 (68.7)	47 (70.1)	46 (68.7)
Response rate $\geq 50\%$	33 (49.3)	41 (61.2)	42 (62.7)
Response rate $\geq 75\%$	15 (22.4)	20 (29.9)	20 (29.9)
Response rate 100%	0	2 (3.0)	3 (6.0)

Abbreviations: CM, chronic migraine; HFEM, high frequency episodic migraine; LOCF, last observation carried forward. Percentages are expressed on column total. HFEM and CM are part of the overall population. Response rate percentages are cumulative.

Table S5. Univariable regressions with baseline selected variables for clinical response (defined as response rate $\geq 50\%$ MHDs) at 3, 6 and 12 months.

Univariable	3 months			6 months			12 months		
	OR	95%CI	p value	OR	95%CI	p value	OR	95%CI	p value
Age (years)	1.00	0.97-1.03	0.92	1.00	0.97-1.03	0.63	1.00	0.97-1.03	0.99
Sex, <i>female</i>	1.65	0.63-4.33	0.30	0.93	0.35-2.49	0.90	1.00	0.37-2.69	0.98
Chronic migraine	0.75	0.25-2.26	0.61	0.94	0.30-2.91	0.92	0.71	0.22-2.30	0.57
Migraine duration (years)	1.00	0.97-1.03	0.62	0.97	0.92-1.01	0.22	0.96	0.91-1.00	0.09
Number of prior failures	0.81	0.61-1.06	0.13	0.92	0.70-1.21	0.55	0.86	0.65-1.14	0.29
Medication overuse	1.62	0.63-4.14	0.30	1.25	0.48-3.23	0.63	1.35	0.52-3.50	0.52
Monthly headache days	0.96	0.91-1.01	0.17	0.96	0.91-1.02	0.24	0.97	0.95-1.03	0.44
Analgesic medication days	0.99	0.95-1.04	0.80	1.00	0.95-1.04	0.99	0.98	0.94-1.03	0.61
Analgesic medication number	0.99	0.98-1.01	0.70	0.98	0.96-1.00	0.09	0.98	0.96-1.00	0.07
MIDAS score	1.00	0.99-1.01	0.71	1.00	0.99-1.01	0.98	0.99	0.99-1.00	0.86
HIT-6 score	0.98	0.91-1.06	0.72	0.95	0.87-1.03	0.26	0.99	0.92-1.07	0.93

Abbreviations: CI confidence interval; CM, chronic migraine; HFEM, high frequency episodic migraine; HIT-6, Headache Impact Test 6; MHDs, monthly headache days; MIDAS, Migraine Disability Assessment; OR, odds-ratio. Patients that discontinued treatment have been considered as non-responders for logistic regressions. For categorical variables, sex male, episodic migraine and absence of medication overuse have been used as reference.