

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Lung Cancer Risks Associated with Occupational Exposure to Pairs of Five Lung Carcinogens: Results from a Pooled Analysis of Case-Control Studies (SYNERGY)

Ann Olsson, Liacine Bouaoun, Joachim Schüz, Roel Vermeulen, Thomas Behrens, Calvin Ge, Hans Kromhout, Jack Siemiatycki, Per Gustavsson, Paolo Boffetta, Benjamin Kendzia, Loredana Radoi, Christine Barul, Stefan Karrasch, Heinz-Erich Wichmann, Dario Consonni, Maria Teresa Landi, Neil E. Caporaso, Franco Merletti, Enrica Migliore, Lorenzo Richiardi, Karl-Heinz Jöckel, Wolfgang Ahrens, Hermann Pohlabein, Guillermo Fernández-Tardón, David Zaridze, John K. Field, Jolanta Lissowska, Beata Świątkowska, John R. McLaughlin, Paul A. Demers, Miriam Schejbalova, Lenka Foretova, Vladimir Janout, Tamás Pándics, Eleonora Fabianova, Dana Mates, Francesco Forastiere, Kurt Straif, Thomas Brüning, Jelle Vlaanderen, and Susan Peters

Table of Contents

Table S1. Pairwise interactions between five occupational lung carcinogens (asbestos, chromium-VI, nickel, PAH, and silica) in relation to lung cancer risk overall and for major lung cancer subtypes in male workers in SYNERGY pooled analyses.

Table S2. Pairwise interactions between five occupational lung carcinogens (asbestos, chromium VI, nickel, PAH, and silica) in relation to lung cancer risk overall and for major lung cancer subtypes in female workers in SYNERGY pooled analyses.

Table S3. Risk of lung cancer (OR), relative excess risk due to interaction (RERI), and test for multiplicative interaction (P-value) associated with single and pairwise combinations of occupational exposure to asbestos, chromium VI, nickel, PAH, and silica in male workers in SYNERGY by smoking status (never, former, and current cigarette smokers).

Figure S1. Heatmaps of Cramer's V statistics (Corr) between the never/ever occupational exposures by lag time in years.