Supplemental Table 1. Data availability across included ECHO cohorts.

| | NICU-HEALTH | ECHO-NOVI | ELGAN | | |
|--|-------------|-----------|------------|--|--|
| Cohort | (N = 380) | (N = 567) | (N = 1089) | | |
| Sample Size (%) with Outcome Variable | | | | | |
| Bayley | 298 (78%) | 438 (77%) | 965 (89%) | | |
| CBCL | 142 (37%) | 550 (97%) | 698 (64%) | | |
| CP | 86 (23%) | 439 (77%) | 952 (87%) | | |
| MCHAT | 196 (51%) | 563 (99%) | 1051 (97%) | | |
| Sample Size (%) with Each Number of Outcome Variable | | | | | |
| 1 | 180 (47%) | 10 (2%) | 46 (4%) | | |
| 2 | 59 (16%) | 109 (19%) | 94 (9%) | | |
| 3 | 141 (37%) | 30 (5%) | 364 (33%) | | |
| 4 | 0 (0%) | 418 (74%) | 585 (54%) | | |

CBCL, Child Behavior Checklist for Ages 1½-5; CP, cerebral palsy; M-CHAT, Modified Checklist for Autism in Toddlers.

Supplemental Table 2. Characteristics of study sample by ECHO cohort

| | NICU-HEALTH | ECHO-NOVI | ELGAN |
|----------------------------------|-------------|-------------|-----------------------------|
| | (N=380) | (N=567) | (N = 1089) |
| Variable | M (SD) or % | M (SD) or % | <i>M</i> (<i>SD</i>) or % |
| Neonatal characteristics | | | |
| Gestational age at birth (weeks) | 26.6 (2.4) | 26.6 (1.9) | 25.6 (1.2) |
| Birth weight (grams) | 952 (427) | 943 (280) | 835 (193) |
| Small for gestational age | 21% | 16% | 13% |
| Male sex | 49% | 54% | 52% |
| Female sex | 51% | 46% | 48% |
| Infant admitted to NICU | 100% | 100% | 100% |
| Infant race | | | |
| American Indian/Alaskan Native | <1% | <1% | <1% |
| Asian | 2.9% | 5.1% | 2.4% |
| Black | 23% | 20% | 27% |
| Multiple Races | 4.2% | 19% | <1% |
| Native Hawaiian/Pacific Islander | <1% | <1% | <1% |
| Other Race | <1% | 7.6% | 7.0% |
| Race Unknown | <2% | <1% | 2.3% |
| White | 68% | 48% | 60% |
| Infant ethnicity | | | |
| Hispanic/Latino/a | 10% | 23% | 12% |
| Not Hispanic/Latino/a | 90% | 77% | 88% |
| • | NICU-HEALTH | ECHO-NOVI | ELGAN |
| | (N = 334) | (N = 498) | (N = 929) |
| | M(SD) or % | M (SD) or % | M(SD) or % |
| Maternal demographics | | | |
| Age at birth (years) | 30.5 (6.1) | 29.1 (6.4) | 28.6 (6.7) |
| 4-year college degree | 34% | 24% | 31% |
| Married/partnered | 56% | 77% | 77% |
| Maternal medical problems | | | |
| Gestational diabetes | 12% | 8.8% | 6.6% |
| Pre-eclampsia | 20% | 25% | 18% |
| Perinatal maternal substance use | | | |
| Tobacco | 5.7% | 16% | 14% |
| Alcohol | 2.6% | 4.0% | 1470 |
| Opioid | 25% | 4.0% 14% | |
| 1 | 25% 3.5% | 9.3% | |
| Marijuana | | | |
| Illicit drugs | 3.1% | 4.6% | |

NICU, neonatal intensive care unit.

No data were available from ELGAN for prenatal alcohol, opioid, marijuana, or illicit drug use.