

Non-coding RNA-related FCGBP downregulation in head and neck squamous cell carcinoma: A novel biomarker for predicting paclitaxel resistance and immunosuppressive microenvironment

Qin Ding^{1,2†}, Fengjie Lin^{1,2†}, Zongwei Huang^{1,2†}, Ying Li^{1,2}, Sunqin Cai^{1,2}, Xin Chen^{1,2}, Hui Liu^{1,2*}, Sufang Qiu^{1,2*}

¹Department of Radiation Oncology, Clinical Oncology School of Fujian Medical University, Fujian Cancer Hospital, Fuzhou, China

²Fujian Provincial Key Laboratory of Translational Cancer Medicine, Fuzhou, China

*Correspondence: Sufang Qiu, sufangqiu@fjmu.edu.cn, Department of Radiation Oncology, Clinical Oncology School of Fujian Medical University, Fujian Cancer Hospital, Fuzhou, 350014, China

Hui Liu, liuhui@fjmu.edu.cn, Department of Radiation Oncology, Clinical Oncology School of Fujian Medical University, Fujian Cancer Hospital, Fuzhou, 350014, China

†These authors have contributed equally to this work.

Supplementary Materials:




Figure S1: Expression analysis and survival analysis for FCGBP in pan-cancer by UALCAN and GEPIA. (A) The expression of FCGBP in pan-cancer based on TCGA cancer and normal data analyzed using UALCAN. (B) The expression of FCGBP in pan-cancer based on TCGA cancer and normal data analyzed using UALCAN using

the GEPIA database. *p value < 0.05; **p value < 0.01; ***p value < 0.001. (C)
Immunohistochemistry results of FCGBP in cancer and normal tissues from the
Human Protein Atlas database.

Table S1: Predicted miRNAs by seven prediction programs.