

Supplemental Table 1. Extended Clinical Information

Cardiogenic Shock Subjects (n=19)

Left Ventricular Ejection Fraction, mean (SD)	24 (15)
% receiving inotropic support	95
% receiving mechanical circulatory support	28

Septic Shock Subjects (n=42)

Norepinephrine-equivalent vasopressor dose, mean (SD)	25.6 (26.1)
---	-------------

Source of Infection, %*:

Pneumonia	57
Urinary Tract	24
Intra-abdominal	21
Skin & Soft Tissue	17
Direct Bloodstream & Endocarditis	7

Organisms, n:

<i>Clostridium difficile</i>	3
Coagulase-negative <i>Staphylococcus</i> species	3
<i>Cornebacterium</i> species	2
<i>Escherichia coli</i>	4
<i>Enterococcus</i> species	4
<i>Haemophilus influenzae</i>	1
Influenza A	3
<i>Klebsiella</i> species	6
Methicillin-resistant <i>Staphylococcus aureus</i>	4
Methicillin-sensitive <i>Staphylococcus aureus</i>	2
<i>Pseudomonas aeruginosa</i>	1
<i>Serratia marcescens</i>	1
<i>Streptococcus agalactiae</i>	1
<i>Streptococcus pneumoniae</i>	2
Unspecified gram negative rods	2
No positive microbiological data	13

*total exceeds 100% as subjects could have multiple sources of infection