

Table S1. Client sampling for client satisfaction survey

		Round 1 (2021)	Round 2 (2022)	Total Clients Surveyed
0 clients	Dispensary	198	212	0
	Health Center	21	15	0
	Hospital	6	2	0
	<b>Total</b>	<b>225</b>	<b>229</b>	<b>0</b>
1 client	Dispensary	55	56	111
	Health Center	10	14	24
	Hospital	0	2	1
	<b>Total</b>	<b>65</b>	<b>72</b>	<b>137</b>
2 clients	Dispensary	164	170	668
	Health Center	30	15	90
	Hospital	8	5	26
	<b>Total</b>	<b>202</b>	<b>190</b>	<b>784</b>
3 clients	Dispensary	0	1	3
	Health Center	0	8	24
	Hospital	0	1	3
	<b>Total</b>	<b>0</b>	<b>10</b>	<b>30</b>
4 clients	Dispensary	0	1	4
	Health Center	0	14	56
	Hospital	0	6	24
	<b>Total</b>	<b>0</b>	<b>21</b>	<b>84</b>
Totals	<b>Total discrete facilities</b>	<b>492</b>	<b>522</b>	
	<b>Total facilities with clients</b>	<b>267</b>	<b>293</b>	
	<b>Total clients surveyed</b>	<b>469</b>	<b>566</b>	<b>1035</b>

**Table S2. Association of health facility type with coverage of Malaria in Pregnancy care components (regression results)**

Health facility type	Malaria in Pregnancy Care Component Coverage*						
	OR [CI] p-value						
	ITN provision	Hb test recorded	mRDT result	Hematinic provision	GA recorded	Any IPTp	IPTp3+
Dispensary	REF	REF	REF	REF	REF	REF	REF
Health Center	<b>0.43</b> [0.32-0.57] <b>p&lt;0.001</b>	<b>1.24</b> [1.11-1.30] <b>p&lt;0.001</b>	0.87 [0.67-1.10] p=0.27	1.03 [0.83-1.27] p=0.79	<b>0.80</b> [0.64-0.99] <b>p=0.04</b>	<b>1.16</b> [1.01-1.32] <b>p=0.03</b>	0.90 [0.80-1.00] p=0.06
Hospital	0.82 [0.43-1.57] p=0.15	<b>1.54</b> [1.25-1.90] <b>p&lt;0.001</b>	0.72 [0.43-1.10] p=0.20	0.74 [0.53-1.05] p=0.08	<b>0.64</b> [0.44-0.91] <b>p=0.01</b>	0.82 [0.63-1.06] p=0.13	<b>1.34</b> [1.08-1.67] <b>p=0.007</b>

Regression analyses include all client records where facility was specified (N=11,256). ITN: insecticide-treated net; Hb: hemoglobin; mRDT: malaria rapid diagnostic test; GA: gestational age; IPTp: intermittent preventive treatment in pregnancy

**Table S3. Association of health facility type with receipt of total number of IPTp doses  
(regression results)**

Health facility type	Total number of IPTp doses received*		
	OR [CI] p-value		
	1	2	3+
Dispensary	REF	REF	REF
Health Center	<b>1.16</b> <b>[1.02-1.33]</b> <b>p=0.025</b>	1.0 [0.87-1.15] p=0.99	0.90 [0.80-1.00] p=0.059
Hospital	0.82 [0.63-1.07] p=0.15	0.78 [0.59-1.02] p=0.07	<b>1.34</b> <b>[1.08-1.67]</b> <b>p=0.007</b>
Unspecified	<b>0.11</b> <b>[0.02-0.84]</b> <b>p=0.033</b>	N/A**	<b>23.13</b> <b>[3.16-169.52]</b> <b>p=0.002</b>

\* Each of the 3 regressions in this table was performed on the subset of women documented as receiving that total number of doses of IPTp. Women who received zero doses (N=2,291) were not included. IPTp: intermittent preventive treatment in pregnancy; \*\*Could not be computed due to small sample size