

SUPPLEMENTARY DIGITAL MATERIAL 1

Supplementary Table I.—Supplementary material, evaluation tools references.

<i>Domain</i>	<i>Evaluation tool</i>	<i>Acronym</i>	<i>References</i>
<i>Clinical features</i>	Markers of complexity	/	Banchero A, Baratto L, Bellelli G, Bellia C, Bernardini B, Bressi M, et al. IPER2: Indicatori di processo e di Esito in Riabilitazione. :76.
	Cumulative Illness Rating Scale	CIRS	Linn BS, Linn MW, Gurel L. CUMULATIVE ILLNESS RATING SCALE. Journal of the American Geriatrics Society. May 1968;16(5):622–6.
	National Institutes of Health Stroke Scale	NIHSS	Goldstein LB, Samsa GP. Reliability of the National Institutes of Health Stroke Scale: Extension to Non-Neurologists in the Context of a Clinical Trial. Stroke. February 1997;28(2):307–10.
	Scale of disability in communication	SDC	Scale of disability in communication Meinecke C, Cappadonia C, Bernardini B. Validazione di una semplice scala di impairment comunicativo nell’anziano. In: XXIII Congresso Nazionale SIMFER Valutazione e qualità dell’assistenza in Medicina Riabilitativa. Ancona; 1995. p. 56–7.
<i>Psycho-social features</i>	Montreal Cognitive Assessment	MoCA	Santangelo G, Siciliano M, Pedone R, Vitale C, Falco F, Bisogno R, et al. Normative data for the Montreal Cognitive Assessment in an Italian population sample. Neurol Sci. April 2015;36(4):585–91.
	Oxford Cognitive Screening	OCS	The Italian OCS Group, Mancuso M, Varalta V, Sardella L, Capitani D, Zoccolotti P, et al. Italian normative data for a stroke specific cognitive screening tool: the Oxford Cognitive Screen (OCS). Neurol Sci. October 2016;37(10):1713–21.
	Activity of Daily Living disabilities: Frenchay Activity Index	FAI	Monteiro M, Maso I, Sasaki AC, Barreto Neto N, Oliveira Filho J, Pinto EB. Validation of the Frenchay activity index on stroke victims. Arq Neuro-Psiquiatr. March 2017;75(3):167–71
	Activity of Daily Living disabilities: modified Functional Walking Categories	mFWC	Perry J, Garrett M, Gronley JK, Mulroy SJ. Classification of Walking Handicap in the Stroke Population. Stroke. June 1995;26(6):982–9.
<i>Functional profile</i>	Modified Barthel Index	mBI	Shah S, Vanclay F, Cooper B. Improving the sensitivity of the Barthel Index for stroke rehabilitation. Journal of Clinical Epidemiology. January 1989;42(8):703–9.
	Trunk Control Test	TCT	Franchignoni FP, Tesio L, Ricupero C, Martino MT. Trunk Control Test as an Early Predictor of Stroke Rehabilitation Outcome. Stroke. July 1997;28(7):1382–5.
	Functional Ambulation Categories	FAC	Geyh S, Kurt T, Brockow T, Cieza A, Ewert T, Omar Z, et al. Identifying the concepts contained in outcome measures of clinical trials on stroke using the international classification of functioning, disability and health as a reference. Journal of Rehabilitation Medicine. 1 August 2004;36(0):56–62.

	Modified Rankin scale	mRS	Quinn TJ, Dawson J, Walters MR, Lees KR. Exploring the Reliability of the Modified Rankin Scale. Stroke. March 2009;40(3):762–6.
	Short Physical Performance Battery	SPPB	Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A Short Physical Performance Battery Assessing Lower Extremity Function: Association With Self-Reported Disability and Prediction of Mortality and Nursing Home Admission. Journal of Gerontology. 1 March 1994;49(2):M85–94.
	Modified Ashworth Scale	MAS	Bohanon RW, Smith MB Interrater reliability of a modified Ashworth scale of muscle spasticity. Phys Ther 1987;67:206-7.
	Fulg-Meyer Assessment scale	FMA	Cecchi F, Carrabba C, Bertolucci F, Castagnoli C, Falsini C, Gnetti B, et al. Transcultural translation and validation of Fugl–Meyer assessment to Italian. Disability and Rehabilitation. 1 May 2020;1–6.