


Supplementary Figure 3 Identification of Enhanced Vaccine Mimotopes for the p15E Murine Cancer Antigen

Shiqi Zhou¹, Yiting Song¹, Yuan Luo¹, Brendan Quinn¹, Yang Jiao¹, Mark D. Long², Scott I. Abrams³, Jonathan F. Lovell^{1*}

¹ Department of Biomedical Engineering, State University of New York at Buffalo, Buffalo, NY 14260, USA

² Department of Biostatistics & Bioinformatics, Roswell Park Comprehensive Cancer Center, Buffalo, NY 14263, USA

³ Department of Immunology, Roswell Park Comprehensive Cancer Center Roswell Park Comprehensive Cancer Center, Buffalo, NY 14263, USA



Supplementary Figure S3. p15E-3C2V immunized mice showed long-term resistance to MC38 tumor (re)challenge. One hundred days after the initial MC38 tumor challenge shown in Figure 5, mice that survived the 3C2V peptide vaccine and 5 new Ctrl mice were inoculated with 1x10⁶ MC38 cells. Tumor volume of **A)** all mice; **B)** one mouse from the wild-type group immunized mouse; **C)** three mice from the 3M group; **D)** two mice from the 3C group; **E)** three mice from the 3C2V group; **F)** one mouse from the 3C6A group; **G)** two mice from the 3C7H group; and **H)** two mice from the 3C7N group. The number above the x-axis indicates the number of tumor-free mice out of all mice from the same group.