


Supplementary Fig.1. Trial sequential analysis for vomiting.



Supplementary Fig.2. Trial sequential analysis for aspiration.

Supplementary Table 1. Results of GRADE evaluation.

Outcome	MD/OR (95%CI)	Number	of	Quality	of	Reasons
		studies		evidence		

Vomiting	0.81(0.58-1.12)	4	$\oplus\oplus\oplus\bigcirc$	Risk of bias
			moderate	was "serious".
Aspiration	0.78(0.45-1.37)	2	$\oplus \oplus \oplus \bigcirc$	Risk of bias
			moderate	was "serious".
Nausea	0.89(0.69-1.15)	5	$\oplus\oplus\oplus\bigcirc$	Risk of bias
			moderate	was "serious".
Degree of	-9.44 (-12.04-	7	$\oplus \oplus \bigcirc\bigcirc$	Risk of bias
thirst	-6.83)		low	was "serious";
				inconsistency
				was "serious".
Anal exhaust	-5.48(-7.74-	5	$\oplus \oplus \bigcirc\bigcirc$	Risk of bias
time	-3.22)		low	was "serious";
				inconsistency
				was "serious".
First	-6.34(-8.90-	3	$\oplus \oplus \bigcirc \bigcirc$	Risk of bias
defecation	-3.79)		low	was "serious";
time				inconsistency
				was "serious".

MD, mean difference; OR, odds ratio; CI, confidence interval