

Supplementary Tables

Supplementary Table 1. Dose Limiting Toxicity Criteria

Toxicity	DLT definition
Hematologic	Grade 3 or febrile neutropenia as per NCI Common Toxicity Criteria for Adverse Events (CTCAE) version 3.0
Allergic reaction/hypersensitivity (including drug fever)	≥Grade 2 as per NCI Common Toxicity Criteria for Adverse Events (CTCAE) version 3.0, and asymptomatic bronchospasm or generalized urticaria
Injection site reaction/extravasation changes	≥ Grade 3 as per NCI Common Toxicity Criteria for Adverse Events (CTCAE) version 3.0
Other non-hematologic	≥Grade 3 as per NCI Common Toxicity Criteria for Adverse Events (CTCAE) version 3.0

Supplementary Table 2. Discontinuation of Dosing in Individual Patients.

→ Use Common Terminology Criteria for Adverse Events (CTCAE) v3.0 unless otherwise specified ←			
CTCAE CATEGORY	ADVERSE EVENT	AGENT	TREATMENT ADJUSTMENT or OTHER ACTION
ALLERGY/ IMMUNOLOGY	≥Grade 2 allergic reaction/hypersensitivity	All drugs used in this study	Discontinue vaccinations and begin observation.
	≥Grade 2 autoimmune reaction (excluding vitiligo)		Discontinue vaccinations and begin observation.
ALL OTHERS	Grade 3 (excluding alopecia)	All drugs used in this study	Discontinue vaccinations and begin observation.